arXiv:1006.5806v1 [hep-th] 30 Jun 2010

TAUP-TH 2913/10
WIS/07/10-JUN-DPPA

Holographic MQCD

Ofer Aharony[a*], David Kutasov[b†], Oleg Lunin[c‡],
Jacob Sonnenschein[d,e§], Shimon Yankielowicz[d,e¶]

[a] *Department of Particle Physics and Astrophysics,*
Weizmann Institute of Science, Rehovot 76100, Israel

[b] *EFI and Department of Physics, University of Chicago, Chicago, IL 60637, USA*

[c] *Department of Physics and Astronomy, University of Kentucky,*
Lexington, KY 40506, USA

[d] *School of Physics and Astronomy,*
The Raymond and Beverly Sackler Faculty of Exact Sciences,
Tel Aviv University, Ramat Aviv, 69978, Israel

[e] *Albert Einstein Minerva Center, Weizmann Institute of Science, Rehovot 76100, Israel*

Abstract

We study a brane configuration of $D4$-branes and $NS5$-branes in weakly coupled type IIA string theory, which describes in a particular limit $d = 4$ $\mathcal{N} = 1$ $SU(N + p)$ supersymmetric QCD with $2N$ flavors and a quartic superpotential. We describe the geometric realization of the supersymmetric vacuum structure of this gauge theory. We focus on the confining vacua of the gauge theory, whose holographic description is given by the MQCD brane configuration in the near-horizon geometry of N $D4$-branes. This description, which gives an embedding of MQCD into a field theory decoupled from gravity, is valid for $1 \ll p \ll N$, in the limit of large five dimensional 't Hooft couplings for the color and flavor groups. We analyze various properties of the theory in this limit, such as the spectrum of mesons, the finite temperature behavior, and the quark-anti-quark potential. We also discuss the same brane configuration on a circle, where it gives a geometric description of the moduli space of the Klebanov-Strassler cascading theory, and some non-supersymmetric generalizations.

July 1, 2010

---

[*] Ofer.Aharony@weizmann.ac.il
[†] dkutasov@uchicago.edu
[‡] lunin.7@gmail.com
[§] cobi@post.tau.ac.il
[¶] shimonya@post.tau.ac.il

1 Introduction and summary

It is believed that in the limit of a large number of colors (*e.g.* large N_c for an $SU(N_c)$ gauge theory), many gauge theories can be reformulated as weakly coupled closed string theories with $g_s \sim 1/N_c$, following the ideas of [1]. This reformulation may facilitate the understanding of non-perturbative properties like confinement and chiral symmetry breaking. For most gauge theories the (higher dimensional) target space of the dual string theory, which is usually referred to as the bulk, is highly curved; so far we do not have good quantitative methods to analyze such string theories. For some specific gauge theories, the dual string theory lives in a weakly curved space. In these cases, the dynamics of the strongly coupled large N_c gauge theory, which lives on a space isomorphic to the boundary of the bulk space, can be studied in detail (see [2] for a review).

Adding N_f flavors in the fundamental representation of an $SU(N_c)$ gauge group corresponds in the dual string theory to adding N_f D-branes in the bulk. For small N_f ($N_f \ll N_c$), the theory with flavor can be studied by adding the dynamics of open strings ending on these D-branes; *e.g.*, when the gauge theory has a global $SU(N_f)$ symmetry, the symmetry currents correspond in the bulk to gauge fields on the D-branes. When N_f becomes of the same order as N_c, the open string coupling on the D-branes, $g_s N_f \sim N_f/N_c$, is not small, and there is no reason to believe that a weakly coupled string description exists.[6]

One way to obtain a weakly coupled string theory for $N_f \sim N_c$ is by gauging the flavor group. In this case the usual arguments of the 't Hooft limit imply that there should be a weakly coupled dual string theory, but it is generally not the same as the original theory, unless the gauge coupling of the flavor group is weak. Of course, when the flavor group is weakly gauged we generally do not expect to get a weakly curved string theory dual, but such a dual may exist when the flavor group is strongly coupled.

In this paper we analyze an example of a large N_c gauge theory with the flavor group gauged, for which a weakly curved string dual exists when the flavor group is strongly coupled. Our large N_c gauge theory will be four dimensional, but we will gauge the flavor group by coupling it to five dimensional gauge fields (with a UV completion given by a six dimensional conformal field theory). The five dimensional flavor gauge theory is IR-free; our weakly curved gravity dual is useful when the interesting physics happens at energies at which this theory is strongly coupled, but there is also a different limit of the same theory where the flavor gauge theory is weakly coupled and the physics is that of the original four dimensional gauge theory with $N_f \sim N_c$ flavors.

Our field theory arises as a decoupling limit of the brane configuration shown in figure 3 below. This configuration (see *e.g.* [4] for a review of the dynamics of this and related brane configurations) involves N $D4$-branes which intersect (along $3+1$ dimensions) two $NS5$-branes with different orientations, and a stack of p additional $D4$-branes which stretch between the fivebranes. This brane system preserves $d = 4$ $\mathcal{N} = 1$ supersymmetry and gives, in a certain decoupling limit, $d = 4$ $\mathcal{N} = 1$ $SU(N+p)$ supersymmetric QCD (SQCD) with $2N$ flavors of "quarks" in the fundamental representation of the gauge group. For vanishing superpotential, this theory flows in the IR to a non-trivial fixed point [8]. The brane construction gives rise to a quartic superpotential which preserves an $SU(N) \times SU(N)$ flavor symmetry. The resulting gauge theory has multiple vacua, some of which are confining (in other vacua, some of the gauge symmetry is spontaneously broken).

---

[6] For a recent review of some approaches to holography for theories with $N_f \sim N_c$, and further references, see [3].

We will discuss a different decoupling limit, in which the flavor symmetry is gauged by coupling it to five dimensional gauge fields (which are the gauge fields on the semi-infinite $D4$-branes in figure 3). We will argue that when $p \ll N$, and the five dimensional gauge theory is strongly coupled (at the characteristic energy scale of the four dimensional dynamics), this theory has a simple string dual. In particular, the confining vacua can be described by placing the MQCD [9] fivebrane in the near-horizon geometry of N $D4$-branes.[7] This gives us a controllable background which is continuously related (by changing parameters) to SQCD, similar to the way that the backgrounds of [5, 6, 7] are continuously related to the pure $d = 4$ $\mathcal{N} = 1$ supersymmetric Yang-Mills (SYM) theory. In our case, there are additional fields living in a higher dimensional space, that only decouple in the limit that their gauge coupling goes to zero. Our main purpose in this paper is to investigate the properties of the resulting system.

We begin in section 2 by describing the brane configuration and its different limits. For $N = 0$, our brane configuration is similar to MQCD, which is obtained by taking the string coupling in figure 3 to be large. We study the brane system in a different limit, where the type IIA string coupling is small, and the MQCD fivebrane is an $NS5$-brane carrying fourbrane flux. To study the dynamics of the fivebrane semiclassically, we take the five dimensional 't Hooft coupling of the $SU(p)$ gauge theory on the $D4$-branes to be large.

For large N, the confining vacua of the gauge theory are described by embedding the MQCD brane configuration (for gauge group $SU(p)$) into the near-horizon geometry of N $D4$-branes. In this sense our discussion provides an embedding of MQCD into a field theory which is decoupled from gravity (the decoupling limit from gravity of the MQCD brane configuration itself is still unknown). Unlike the original MQCD configuration, this enables us to have normalizable states (corresponding to four dimensional particles) coming from the MQCD brane, and to study the theory at finite temperature. We discuss both the brane configuration corresponding to the SQCD theory discussed above, and its generalization to the case where the $D4$-branes live on a circle; in the latter case our field theories are $SU(N) \times SU(N+p)$ gauge theories similar to the ones that appear in the Klebanov-Strassler cascade [6], but we study these theories in a different range of parameters from [6]. As in the non-compact case, we find that some features of the cascading theories (like their moduli space) are realized in our limit as well, while other features are different.

In section 3 we analyze the field theories corresponding to our brane configurations, and in particular study their moduli space and match it to the dual string description, finding precise agreement whenever the string theory computation is under control. In particular, we find an elegant geometrical description for the complicated moduli space [11] of the Klebanov-Strassler cascading theory. In section 4 we compute the spectrum of operators and states in our string theory dual. We find that the spectrum is generally continuous from the four dimensional point of view, because some higher dimensional fields do not decouple in our limit, but there are also some discrete states which may be continuously connected to the mesons of SQCD.

---

[7] Holography for the $D4$-brane geometry was developed in [10].

In section 5 we discuss some of the energy scales in our problem, and in particular the quark-anti-quark potential. We show that for some range of parameters this is dominated by the five dimensional IR-free physics, but that there is also a range of parameters (and of quark-anti-quark distances) for which it is dominated by the four dimensional confining physics. In section 6 we discuss the behavior of our system at finite temperature, showing that at all finite temperatures the confining phase has a higher free energy than the Higgs phase, in agreement with field theory expectations. Finally, in section 7 we discuss some non-supersymmetric generalizations of our construction, including a case where there is a first order finite temperature phase transition (at which the fivebrane falls into the horizon). Clearly there are many possible generalizations of our setup, both supersymmetric (*e.g.* theories related to $d = 3$ $\mathcal{N} = 2$ SQCD) and non-supersymmetric; we leave their analysis to future work.

2 The brane construction

2.1 $\mathcal{N} = 1$ Supersymmetric Yang-Mills from type IIA string theory

Pure $d = 4$ $\mathcal{N} = 1$ SYM with gauge group $U(p)$ can be realized in type IIA string theory as the low energy limit of the system of intersecting $D4$-branes and $NS5$-branes depicted in figure 1. All the branes in the figure are extended in the $\mathbb{R}^{3,1}$ labeled by (x_0, x_1, x_2, x_3). The fivebranes are further extended in

$$\begin{aligned} NS: &\qquad v = x_4 + ix_5, \\ NS': &\qquad w = x_8 + ix_9, \end{aligned} \tag{1}$$

while the p fourbranes form a line segment of length L in the x_6 direction,

$$-\frac{L}{2} \leq x_6 \leq \frac{L}{2}. \tag{2}$$

As reviewed in [4], this brane configuration preserves $\mathcal{N} = 1$ supersymmetry in the $3 + 1$ dimensions common to all the branes, (0123). All the fields in the $D4$-brane gauge theory other than the $U(p)$ gauge fields and gauginos get masses of order $1/L$ due to the boundary conditions at $x_6 = \pm L/2$.



Figure 1: The brane system realizing $d = 4$ $\mathcal{N} = 1$ SYM with gauge group $U(p)$.

The classical four dimensional gauge coupling is given by

$$g_{YM}^2 = \frac{g_s l_s}{L}. \tag{3}$$

Later, we will be interested in the large p limit of this system, in which the strength of interactions is governed by the 't Hooft coupling,

$$\lambda_p^{(4)} = g_{YM}^2 p = \frac{\lambda_p}{L}; \qquad \lambda_p \equiv g_s l_s p. \tag{4}$$

λ_p (which has units of length) is the $4+1$ dimensional 't Hooft coupling of the $D4$-brane theory.

Quantum mechanically, the coupling of the four dimensional gauge theory runs with the scale. One can view $1/L$ as the UV cutoff of this theory, and $\lambda_p^{(4)}$ as the value of the coupling at the UV cutoff scale. In order for the dynamics of the brane configuration to reduce to that of $\mathcal{N} = 1$ SYM at energies well below the cutoff scale, this coupling must be taken to be very small, $\lambda_p^{(4)} \ll 1$.

The classical picture of $D4$-branes ending on $NS5$-branes (figure 1) is qualitatively modified by g_s effects [12]. To exhibit these effects, it is convenient to view type IIA string theory at finite g_s as M-theory compactified on a circle of radius $R = g_s l_s$. Both the $D4$-branes and the $NS5$-branes correspond from the eleven dimensional point of view to $M5$-branes, either wrapping the M-theory circle ($D4$-branes), or localized on it ($NS5$-branes). The configuration of figure 1 lifts to a single $M5$-brane, which wraps $\mathbb{R}^{3,1}$ and a two dimensional surface in the $\mathbb{R}^5 \times S^1$ labeled by (v, w, z). Here

$$z \equiv x_6 + i x_{11}, \tag{5}$$

and x_{11} parameterizes the M-theory circle, $x_{11} \sim x_{11} + 2\pi R$. The shape of the fivebrane in $\mathbb{R}^5 \times S^1$ is described by the equations [9]

$$vw = \xi^2, \qquad v = \xi e^{-z/pR} = \xi e^{-z/\lambda_p}, \tag{6}$$

where without loss of generality we can choose ξ to be real and positive. Note that:

- The classical brane configuration of figure 1 lies on the surface $vw = 0$, while the quantum shape is deformed away from this surface. In particular, the $D4$-branes, that are classically at $v = w = 0$, are replaced in the quantum theory by a tube of width $\sim \xi$ connecting the (deformed) NS and NS'-branes. Indeed, defining the radial coordinate u via

  $$u^2 = |v|^2 + |w|^2, \tag{7}$$

  we see that (6) satisfies $u \geq \sqrt{2}\xi$. One can interpret ξ/l_s^2 as the brane analog of the dynamically generated scale of the $\mathcal{N} = 1$ SYM theory.

- Classically, the brane configuration is confined to the interval (2), while quantum mechanically it extends to arbitrarily large $|x_6|$. For example, for large positive x_6, the fivebrane takes the shape

  $$z \simeq \lambda_p \ln(w/\xi); \quad v \simeq 0. \tag{8}$$

  One can think of (8) as describing an NS'-brane deformed by the $D4$-branes ending on it from the left. Since these $D4$-branes are codimension two objects on the fivebrane, they give rise to a deformation of it that does not go to zero at infinity.

- As usual, isometries in the bulk give rise to global symmetries of the field theory on the branes. Consider the following three $U(1)$ symmetries: $U(1)_{45}$, corresponding to rotations in the (45) plane, $U(1)_{89}$, corresponding to rotations in the (89) plane, and $U(1)_{11}$, corresponding to translations in x_{11}. Classically, the first two symmetries are preserved by the brane configuration of figure 1, while the third one is broken by the positions of the $NS5$-branes. Quantum mechanically, the asymptotic $x_6 \to \infty$ shape (8) breaks one linear combination of $U(1)_{89}$ and $U(1)_{11}$, while its analog as $x_6 \to -\infty$ breaks a linear combination of $U(1)_{45}$ and $U(1)_{11}$. The full brane configuration (6) preserves a single $U(1)$ symmetry, whose action is given by

  $$v \to e^{i\alpha} v, \qquad w \to e^{-i\alpha} w, \qquad x_{11} \to x_{11} - \alpha\lambda_p. \tag{9}$$

  Note that the other two $U(1)$ symmetries are broken by the asymptotic boundary conditions, so they are broken explicitly from the point of view of the field theory living on the branes. As discussed in [9], since pure $\mathcal{N} = 1$ SYM theory has no unbroken $U(1)$ global symmetry, all states that are charged under the symmetry (9) are expected to decouple in any limit that leaves only the degrees of freedom of this four dimensional gauge theory. One of the broken $U(1)$ symmetries discussed above is actually not completely broken by the asymptotic shape (8) and its $x_6 \to -\infty$ analog. It is broken to Z_p, which is further spontaneously broken by the full brane configuration (6). This $U(1)$ symmetry can be identified with the (anomalous) R-symmetry of the four dimensional SYM theory.

- At large u, the distance between the NS and NS'-branes goes to infinity; hence the separation L appears to be ill defined. This is not surprising, since (3) relates L to the four dimensional gauge coupling, which changes with the scale. To define it, one can take the radial coordinate u (7) to be bounded, $u \leq u_\infty$, and demand that at $u = u_\infty$, $x_6 = \pm L/2$. Assuming that the four dimensional 't Hooft coupling $\lambda_p^{(4)}$ (4) is very small, this gives the following relation among the different scales:

  $$\xi = u_\infty \exp\left(-L/2\lambda_p\right) = u_\infty \exp\left(-1/2\lambda_p^{(4)}\right). \tag{10}$$

  This is the brane analog of the relation between the QCD scale and the gauge coupling in SYM theory, with u_∞/l_s^2 playing the role of a UV cutoff.[8] As in SYM, we can remove the cutoff, by sending $u_\infty, L/\lambda_p \to \infty$ while keeping the "QCD scale" ξ/l_s^2 fixed.

- Although the curved fivebrane is non-compact, and in particular extends to infinity in $\mathbb{R}^5$, the non-trivial dynamics is restricted to the intersection region. The only low energy modes that live on the fivebrane at large (v, w, x_6) are $5+1$ dimensional free fields that describe the position of a single fivebrane, and their superpartners. One can take a limit in which these fields decouple, and only the $3+1$ dimensional physics remains, but we will not do that here.

As one varies the parameters of the brane system of figure 1, the language in terms of which its IR dynamics is most usefully described changes. The pure SYM description is valid when $\lambda_p^{(4)}$ (4) and the dynamically generated scale ξ (6) are small. In this regime the brane description reduces to the gauge theory one.

In other regions in parameter space the dynamics can be studied by analyzing the geometry of the branes. One such region is obtained by sending $g_s \to \infty$, *i.e.* taking the radius of the M-theory circle, R, to be much larger than the eleven dimensional Planck length. If ξ is also taken to be sufficiently large, (6) describes a large and smooth $M5$-brane, whose dynamics can be studied semiclassically. The low energy theory of this fivebrane is known as MQCD; it was discussed in [9] and subsequent work. Since it is related to $\mathcal{N} = 1$ SYM by a continuous deformation of the parameters of the brane configuration, and no phase transitions are expected along this deformation, the two theories are believed to be in the same universality class. However, many of their detailed features are expected to be different.

In this paper we will focus on a different region in the parameter space of the brane system. We will take the type IIA string coupling to be small, but p to be large, such that the five dimensional 't Hooft coupling is large, $\lambda_p \gg l_s$. As mentioned above, for large L and small ξ the low energy dynamics of this system reduces to $\mathcal{N} = 1$ SYM. On the other hand, if ξ is sufficiently large, the fivebrane described by (6) is weakly curved (in string units). Thus, the situation is similar to that in MQCD, except for the fact that the string coupling is weak. This implies that the fivebrane in question is an $NS5$-brane, which also carries RR six-form flux ($D4$-brane charge).

---

[8] The corresponding energy scale is in general different from the KK scale $1/L$ mentioned above.

The shape of this $NS5$-brane is given by the reduction of the eleven dimensional profile (6) to ten dimensions. It is parameterized by two functions of x_6, u and α, which are defined (on the fivebrane) by

$$v = ue^{i\phi}\cos(\alpha), \qquad w = ue^{-i\phi}\sin(\alpha). \tag{11}$$

Looking back at (6), we see that u and α are given by

$$u = \xi\sqrt{2\cosh\left(\frac{2x_6}{\lambda_p}\right)}, \qquad \tan(\alpha) = \exp\left(2x_6/\lambda_p\right). \tag{12}$$

The configuration (6) also involves a non-zero expectation value for the (compact) scalar field which labels the position of the $NS5$-brane along the M-theory circle. As ϕ in (11) varies between 0 and 2π, the scalar field winds p times around the circle, giving the $NS5$-brane its $D4$-brane charge.

When $\lambda_p, \xi \gg l_s$, the shape (12) is weakly curved and a semiclassical description should be reliable. This description depends on whether the back-reaction of the branes on the geometry can be neglected. The p $D4$-branes modify the geometry around them significantly up to distances of order $(\lambda_p l_s^2)^{1/3}$. Thus, if ξ satisfies the constraint

$$\xi^3 \gg \lambda_p l_s^2, \tag{13}$$

one can neglect the back-reaction and treat the curved fivebrane as a probe in flat spacetime. Otherwise, the back-reaction is important, and one can try to replace the branes by their geometry, and describe their low-energy dynamics using holography. This is an interesting problem that we will leave to future work.

The regime (13), where the curved fivebrane (11), (12) can be treated as a probe, is quite analogous to MQCD. The validity of the semiclassical analysis of the fivebrane relies in this case on the 't Hooft large p limit rather than on large g_s, as in [9], but many of the qualitative properties are similar.

2.2 A holographic embedding of MQCD

We next embed the brane configuration of figure 1 in a larger system, which is more amenable to a holographic analysis. To do this, we add N infinite $D4$-branes stretched in the x_6 direction (see figure 2). These branes do not break any of the supersymmetries preserved by the configuration of figure 1. Thus, we can place them anywhere in the $\mathbb{R}^5$ labeled by (v, w, x_7), without influencing the shape of the curved $NS5$-brane (12). This can be seen directly by replacing the N $D4$-branes by their geometry, and studying the dynamics of the curved $NS5$-brane (12) in that geometry. This description should be valid for $N \gg p$, and we will restrict to this ("probe") regime below.



Figure 2: The brane system with N additional infinite $D4$-branes.

Viewing the N $D4$-branes as $M5$-branes wrapped around the M-theory circle, their eleven dimensional geometry is given by[9]

$$\begin{aligned} ds^2 &= H^{-1/3}\left(dx_\mu^2+dx_6^2+dx_{11}^2\right)+H^{2/3}\left(|dv|^2+|dw|^2+dx_7^2\right), \\ C_6 &= H^{-1}d^4x\wedge dx_6\wedge dx_{11}, \qquad H=1+\frac{\pi\lambda_N l_s^2}{|\vec{r}-\vec{r}_0|^3}, \end{aligned} \tag{14}$$

where $\lambda_N = g_s l_s N$ is the $4+1$ dimensional 't Hooft coupling of the N $D4$-branes (defined as in (4)), $\mu = 0,1,2,3$, $\vec{r}=(v,w,x_7)$ labels position in $\mathbb{R}^5$, with $\vec{r}=0$ corresponding to the classical position of the p $D4$-branes in figure 2, and $\vec{r}_0\in\mathbb{R}^5$ is the position of the N $D4$-branes.

For $g_s\neq 0$, the $NS5$-branes and p $D4$-branes form a curved $M5$-brane whose shape may be obtained by plugging the ansatz

$$v=u(x_6)e^{i\phi(x_{11})}\cos(\alpha(x_6)), \qquad w=u(x_6)e^{-i\phi(x_{11})}\sin(\alpha(x_6)), \tag{15}$$

into the fivebrane worldvolume action. Parametrizing the $M5$-brane worldvolume by the coordinates (x_μ, x_6, x_{11}), the induced metric corresponding to (15) takes the form

$$ds_{ind}^2 = H^{-1/3}\left\{dx_\mu^2+\left[1+H\left((u\alpha')^2+(u')^2\right)\right]dx_6^2+\left(1+H(u\dot{\phi})^2\right)dx_{11}^2\right\}, \tag{16}$$

where $u'\equiv\partial_{x_6}u$, $\alpha'\equiv\partial_{x_6}\alpha$ and $\dot{\phi}\equiv\partial_{x_{11}}\phi$. The Lagrangian is[10]

$$L = H^{-1}\sqrt{1+H(u\dot{\phi})^2}\sqrt{1+H((u\alpha')^2+(u')^2)}-H^{-1}. \tag{17}$$

---

[9] It is well known that dimensional reduction of this geometry gives the correct description of $D4$-branes in type IIA string theory. Thus, the discussion below is valid in the weakly coupled type IIA limit as well.

[10] Here and below, we omit the tension of the fivebrane, which appears as a multiplicative factor in the Lagrangian.

The equations of motion imply that $\dot{\phi}$ must be constant; thus $\phi = x_{11}/pR = x_{11}/\lambda_p$. The Noether charges J, E associated with the invariances under the shifts of α and x_6, respectively, are then given by

$$\begin{aligned} J &= \frac{u^2\alpha'\sqrt{1+Hu^2/\lambda_p^2}}{\sqrt{1+H\left[(u\alpha')^2+(u')^2\right]}}, \\ E &= H^{-1} - \frac{H^{-1}\sqrt{1+Hu^2/\lambda_p^2}}{\sqrt{1+H\left[(u\alpha')^2+(u')^2\right]}}. \end{aligned} \tag{18}$$

Supersymmetric configurations have $E = 0$. Substituting into (18) we find:

$$\begin{aligned} \alpha' &= J/u^2, \\ (u')^2 &= \frac{u^2}{\lambda_p^2} - \frac{J^2}{u^2}. \end{aligned} \tag{19}$$

Note that the equations (19) that determine the shape of the supersymmetric $M5$-brane are independent of the form of the harmonic function H, and in particular of the positions of the N $D4$-branes in figure 2. This agrees with the expectation that there is no force between the various branes. As a check, one can verify that the profile (12) indeed solves (19), with

$$J\lambda_p = 2\xi^2. \tag{20}$$

In this solution α goes from $\alpha(x_6 \to -\infty) = 0$ to $\alpha(x_6 \to \infty) = \pi/2$.

So far, our discussion took place in the full type IIA string theory. We next take a decoupling limit, by omitting the 1 in the harmonic function H (14). This corresponds to studying the brane configuration of figure 2 in the $(2,0)$ theory of N $M5$-branes, compactified on a circle of radius R. One can think of the curved $M5$-brane (12) as a localized defect in this theory. The dynamics of the modes of the $(2,0)$ theory contributes to the interactions among the fields localized on the defect, which include four dimensional $\mathcal{N} = 1$ gauge superfields. We will see later that the low energy theory contains some additional higher dimensional modes, and is thus not purely $\mathcal{N} = 1$ SYM. However, it does not contain any gravitational or stringy dynamics, in contrast to MQCD and its weakly coupled analog described in the previous subsection.

As mentioned above, the shape of the curved $M5$-brane (12) is independent of the position of the N $D4$-branes, $\vec{r}_0$. Consider the limit $\vec{r}_0 \to 0$, in which all the $D4$-branes in the classical configuration of figure 2 are coincident (see figure 3). As is clear from the figure, there are states (corresponding to fundamental strings stretched between the two stacks of $D4$-branes), whose masses go to zero in this limit. We will discuss their field theoretic interpretation in the next section; here we note that for $\vec{r}_0 = 0$ the system has some additional supersymmetric vacua.

In the classical brane description, these vacua can be obtained as follows. One or more of the fourbranes stretched between the $NS5$-branes can connect to semi-infinite branes



Figure 3: The brane system of figure 2 at the origin of its moduli space.

attached from the left to the NS-brane, and make semi-infinite $D4$-branes stretching from the NS'-brane to $x_6 = -\infty$. As is clear from figure 3, this leaves behind semi-infinite fourbranes stretching from the NS-brane to $x_6 = +\infty$. The semi-infinite branes in the two stacks can now be independently displaced along the appropriate $NS5$-brane, thus giving rise to a new branch of the moduli space of brane configurations. In this branch, the number of $D4$-branes stretching between the $NS5$-branes is reduced.

When all p $D4$-branes reconnect as described above, the fivebrane splits into two separate fivebranes, each of which corresponds to a solution of (19) with $J = 0$. The angle α (11) takes the constant values 0 and $\pi/2$ for the NS and NS'-brane, respectively. The solution to (19) is

$$u = K \exp\left(\pm x_6/\lambda_p\right). \tag{21}$$

The number of $D4$-branes stretched all the way from $x_6 = -\infty$ to $+\infty$ (*i.e.* the background $D4$-brane flux) decreases to $N - p$ (see figure 4). We will discuss the field theory interpretation of these extra branches of the moduli space of brane configurations in the next section.

Before introducing the N extra $D4$-branes, we found that the parameter ξ, (6), (12), must satisfy the bound (13) in order for the back-reaction of the p $D4$-branes on the geometry to be negligible. In the near-horizon geometry of the N $D4$-branes, the back-reaction of the p $D4$-branes is always a subleading effect in p/N (which we are assuming is small). There is still a constraint coming from the fact that if ξ is small, the tube connecting the $NS5$-branes is located in the region where the curvature of the near-horizon geometry is large and cannot be trusted. For $\vec{r}_0 = 0$ this constraint takes the form

$$\frac{\xi}{l_s^2} \gg \frac{1}{\lambda_N}. \tag{22}$$

It is much less stringent than (13); we will impose it below. Recall that ξ/l_s^2 is the QCD scale associated with the tube (6); it is held fixed in the decoupling limit from



Figure 4: The "disconnected" branch of the moduli space.

gravity described above. The constraint (22) is precisely the requirement that the $4+1$-dimensional $U(N)$ gauge theory on the $D4$-branes is strongly coupled at the scale ξ/l_s^2 [10].

2.3 Compact x_6

Another interesting brane system is obtained from that of figure 2 by compactifying the x_6 direction on a circle, $x_6 \sim x_6 + 2\pi R_6$. The resulting configuration (with $\vec{r}_0 = 0$) is depicted in figure 5. While the bosons living on the branes must satisfy periodic boundary conditions around the circle, fermions can be either periodic or antiperiodic. Most of our discussion below will involve the periodic case, in which supersymmetry is unbroken by the boundary conditions. In this case, the string background is still given by (14), and the shape of the curved fivebrane by (11), (12), (21), with x_6 periodically identified.

On the cylinder labeled by (x_6, u), the fivebrane described by (11), (12) takes the form depicted in figure 6. Starting at the boundary $u = u_\infty$, it spirals down the cylinder and then climbs back up to the boundary. In the process, the angle α changes by $\pi/2$. The disconnected configuration (21) gives rise to a solution with constant α, which spirals down the cylinder and does not climb back.

Although in the covering space of the cylinder the shape of the fivebrane is the same as in the non-compact case, there are two important differences between the two cases. The first has to do with the fact that the curved fivebrane carries p units of fourbrane charge. In the non-compact case, if one fixes x_6 and increases the radial coordinate u, the flux increases by p units, from N to $N+p$, when one crosses the fivebrane. For finite R_6, increasing u at fixed x_6 one encounters the fivebrane multiple times, as it spirals around the cylinder. At each encounter, the fourbrane flux increases by p units. If we fix the flux to be $N+p$ at the UV boundary $u = u_\infty$, as we spiral down, it may[11] eventually

---

[11] The change of flux between $u \sim \xi$ and u_∞ is $\Delta N \sim L/R_6$, where L is the distance between the $NS5$-branes (in the covering space) evaluated at u_∞ (see the discussion around (10)). It could be large or small compared to N.



Figure 5: N $D4$-branes that wrap the x_6 circle, and p that are stretched between the NS and NS'-branes.

become comparable to p, where the probe approximation breaks down.

The second difference between the compact and non-compact cases concerns the vacuum structure of the model. In the non-compact case, fivebranes described by (11), (12) with different values of ξ correspond to different theories, much like SYM theories with different values of Λ_{QCD}. Indeed, for large u one has (see (8))

$$x_6 \sim \lambda_p \ln\left(\frac{u}{\xi}\right) , \tag{23}$$

and changing ξ corresponds to a non-normalizable mode in the geometry (14). In the compact case, profiles with

$$\xi = \xi_n = \xi_0 e^{\frac{2\pi n R_6}{\lambda_p}} , \qquad n \in Z \tag{24}$$

give rise to the same asymptotics (23), and thus describe different vacua of a single theory. The integer n is bounded from above by the requirement $\xi_n < u_\infty$ and from below by the breakdown of the probe approximation described above. Later we will discuss the field theory interpretation of these vacua.

When the radius R_6 becomes too small, the adjacent coils in figure 6 approach each other, and one can no longer ignore their back reaction on the geometry. The requirement that this does not happen leads to the inequality

$$R_6 \gg \left(\frac{\lambda_p^4}{\lambda_N}\right)^{1/3} . \tag{25}$$

Another constraint on R_6 comes from requiring that upon reducing the geometry (14) to IIA string theory, the radius of the x_6 circle at $u \sim \xi$ should be much larger than the string length. This leads to the constraint

$$R_6 \gg \left(\frac{\lambda_N l_s^2}{(\xi/l_s^2)^3} \right)^{1/6} . \tag{26}$$



Figure 6: For compact x_6, the curved $NS5$-brane spirals down the (x_6, u) cylinder and then climbs back up. The downward (upward) part of the spiral is colored red (blue).

The brane configuration of figure 5 is T-dual to the system studied in [6], which involves regular and fractional $D3$-branes placed at a conifold singularity in type IIB string theory. The Klebanov-Strassler theory describes the limit of our configuration as $R_6 \to 0$ and $\lambda_p, \lambda_N \to 0$, keeping the four dimensional QCD scale ξ/l_s^2 fixed; this is an opposite limit to the limit of strong five dimensional gauge coupling that we focus on in this paper. On the type IIB side the brane sources are replaced by varying three-form and five-form fluxes. In particular, the gauge-invariant charge associated with F_5 changes along the throat; this change can be attributed to a sequence of Seiberg dualities (see section 3). One can also define a Page charge associated with the five-form [13], which is conserved along the flow, but is not gauge-invariant (though its value modulo p is gauge-invariant).

The type IIA story is simpler. In the probe approximation ($p \ll N$) discussed in this paper, the four-form field strength has explicit sources (the spiraling fivebrane), and there is only one charge, which is both gauge-invariant and conserved. It changes as one moves in the radial direction due to the presence of sources.

The system discussed in [6] is known to have a rich vacuum structure, some of which is described by regular type IIB supergravity solutions (in a certain regime in the parameter space of brane configurations). The type IIA description in terms of a fivebrane with

$D4$-brane charge winding around the cylinder in figure 6 is valid in a different regime in parameter space, but the (supersymmetric) vacuum structure is expected to be the same. We will comment on this comparison further below.

3 Field theory

In this section we will discuss the field theory that governs the low energy dynamics of the brane configurations of figures 3 and 5, and its relation to that of the brane systems described above.

3.1 Non-compact x_6

We start with the brane configuration of figure 3, and consider its low energy dynamics in the weak coupling limit. The low energy effective field theory on the $D4$-branes contains two types of excitations: four dimensional fields localized in the intersection region, and five dimensional fields living on the semi-infinite fourbranes at $|x_6| \geq L/2$.

The four dimensional fields include a $U(N+p)$ $\mathcal{N}=1$ SYM theory, and two sets of N (anti) fundamental chiral superfields $(Q^i_L, \tilde{Q}_{L,i})$, $(Q^\alpha_R, \tilde{Q}_{R,\alpha})$; $i, \alpha = 1, 2, \cdots, N$. The latter are coupled by the superpotential [14]

$$W_\lambda = \lambda(\tilde{Q}_L \cdot Q_R)(\tilde{Q}_R \cdot Q_L) \ , \tag{27}$$

where the scalar product stands for contraction of the $U(N+p)$ color indices, and the flavor indices (i, α) are contracted between the two gauge-invariant bilinears. This theory is invariant under a $U(N)_L \times U(N)_R$ global symmetry, which acts on the indices (i, α) (note that this is a non-chiral symmetry). This symmetry is the global part of the gauge group of the semi-infinite $D4$-branes on the left and right of figure 3.

The theory is also invariant under a $U(1)$ global symmetry under which Q_L, $\tilde{Q}_L$ have charge one, and Q_R, $\tilde{Q}_R$ have charge minus one. This symmetry corresponds in the brane picture to the $U(1)$ symmetry (9). To see that, it is useful to think about it as a difference of two R-symmetries, acting on $(Q_L, \tilde{Q}_L)$ and $(Q_R, \tilde{Q}_R)$, respectively. These R-symmetries correspond in the brane language to rotations in the w and v planes. Hence, their difference, which does not act on the supercharges, acts on v and w as in (9).

While the chiral superfields are localized in x_6 (at $x_6 = \pm L/2$), the vector superfields are five dimensional fields living on the line segment (2). At energies of order $1/L$, one starts seeing the massive Kaluza-Klein (KK) states, both of the gauge field and of the other $D4$-brane modes (the five transverse scalars and fermions). The four dimensional gauge coupling is given in terms of the five dimensional one by an analog of equation (4), $\lambda^{(4)}_{N+p} = \lambda_{N+p}/L$. In this section we assume that it is small at the KK scale, though this is not true when the gravitational approximation of the holographic description of the previous section is valid.

In addition to the fields mentioned above, the brane system contains five dimensional fields living on the two stacks of semi-infinite fourbranes in figure 3. They are described by a $4+1$ dimensional SYM theory with sixteen supercharges, broken down to eight supercharges by the boundary conditions at $x_6 = \pm L/2$. The supersymmetry is further broken down to $\mathcal{N} = 1$ by the couplings of the five dimensional fields to the four dimensional ones. These couplings can be read off from figure 3 by examining the effects of geometric deformations. In particular, the superpotential (27) receives an additional contribution of the form

$$W_\phi = \lambda_\phi \left(\tilde{Q}_L \cdot Q_L \Phi_L + \tilde{Q}_R \cdot Q_R \Phi_R \right) , \tag{28}$$

for some constant λ_ϕ, where Φ_L is an $N \times N$ matrix chiral superfield that parametrizes the position of the left semi-infinite $D4$-branes along the NS-brane (*i.e.* in the v direction). This field is defined on the half-infinite line segment $x_6 \leq -L/2$, but what enters W_ϕ is only its value at $x_6 = -L/2$. Similarly, Φ_R parametrizes the position in w of the semi-infinite $D4$-branes on the right of figure 3, $x_6 \geq L/2$, and what enters (28) is its value at $x_6 = L/2$.

As mentioned above, the $U(N) \times U(N)$ global symmetry of the four dimensional theory is part of the gauge symmetry of the five dimensional theory. Similarly, the vacuum expectation values of the five dimensional fields Φ_L and Φ_R correspond from the four dimensional point of view to parameters in the Lagrangian – turning them on gives masses to $(Q_L, \tilde{Q}_L)$ and $(Q_R, \tilde{Q}_R)$, respectively.

The 't Hooft coupling of the five dimensional theory, λ_N (see (4)), has units of length. In section 2 we discussed the strong coupling regime, in which this length is much larger than l_s. The gauge theory description is valid in the opposite limit, $\lambda_N \ll l_s$. As usual, many aspects of the supersymmetric vacuum structure are insensitive to the coupling, and can be compared between the two regimes.

Varying the superpotential $W = W_\lambda + W_\phi$ with respect to $\tilde{Q}_L$ gives the F-term equation

$$\lambda Q_R \tilde{Q}_R \cdot Q_L + \lambda_\phi Q_L \Phi_L = 0 \ . \tag{29}$$

Three similar equations are obtained by varying with respect to the other components of Q, $\tilde{Q}$. Some of the solutions of (29) and of the D-term constraints can be described as follows.

One branch of solutions is $Q_{L,R} = \tilde{Q}_{L,R} = 0$, with Φ_L, Φ_R general diagonal matrices. The corresponding brane configuration is given in figure 7. As is clear both from the field theory analysis and from the brane perspective, in this branch the chiral superfields are massive, and the theory generically reduces at low energies to $\mathcal{N} = 1$ pure SYM with gauge group $U(N+p)$. This takes us back to the discussion of section 2.1.

Another branch of supersymmetric solutions corresponds to the brane configuration of figure 4. At strong coupling, the resulting vacua were discussed in section 2, around equation (21). At weak coupling, they can be described as follows.

In order for the $(N-p)$ $D4$-branes stretched from $-\infty$ to ∞ in figure 4 not to intersect the $NS5$-branes, we must displace them in (v, w) and/or x_7. In the gauge theory, the former corresponds to turning on non-zero expectation values for the five dimensional



Figure 7: The brane realization of vacua with $\Phi_L, \Phi_R \neq 0$.

fields Φ_L, Φ_R. Setting these expectation values to zero and displacing the fourbranes in x_7 corresponds to turning on Fayet-Iliopoulos (FI) D-terms for the diagonal $U(1)$'s in $U(N-p)$ subgroups of the two $U(N)$'s mentioned above. Denoting these D-terms by $x_7^{L,R}$, the D-term potential sets

$$\begin{aligned} Q_L^\dagger \cdot Q_L &= x_7^L I_{N-p} \ , \\ Q_R^\dagger \cdot Q_R &= x_7^R I_{N-p} \ , \end{aligned} \tag{30}$$

where we assumed that $x_7^{L,R}$ are positive, and set $\tilde{Q}_{L,R} = 0$ for the $(N-p)$ flavors involved in (30). The latter is necessary for the F-term conditions (29) to be satisfied. Indeed, since we set $\Phi_L = 0$ for the $(N-p)$ flavors in (30), the F-term constraint requires $\tilde{Q}_R \cdot Q_L$ to vanish. Since Q_L is non-singular in a $(N-p) \times (N-p)$ block constrained by (30), $\tilde{Q}_R$ must vanish in that block. To solve (30), we take $Q_L = \sqrt{x_7^L}\delta_A^i$, where $i = 1, \cdots, N-p$ is a flavor index, and $A = 1, \cdots, N-p$ a color one. Similarly, we take $Q_R = \sqrt{x_7^R}\delta_A^\alpha$, where α is the flavor index, and the color index A runs over the same range as for Q_L. The D-term potential in the $U(N+p)$ four dimensional gauge theory, which requires $Q_L^\dagger Q_L = Q_R^\dagger Q_R$, then implies that one must have $x_7^L = x_7^R$, a constraint that is obvious from the geometry of figure 4.

The above discussion takes care of $(N-p)$ colors and flavors. This leaves $2p$ colors and p flavors of Q, $\tilde{Q}$. To solve the D and F-term constraints, we take Q_L, $\tilde{Q}_L$ to be diagonal and non-zero in a $p \times p$ block with $i, A = N-p+1, \cdots, N$, and Q_R, $\tilde{Q}_R$ to be diagonal and non-zero in a $p \times p$ block with $\alpha, A = N+1, \cdots, N+p$. The F-term constraints (29) allow us to turn on an arbitrary Φ_L in the flavor sector with non-zero Q_R and vice-versa. Of course, the above forms of Q, $\tilde{Q}$ are up to gauge and global transformations.

The above discussion involved vacua of the low energy gauge theory in which the four dimensional $U(N+p)$ gauge group is completely broken. There are other branches of the moduli space of supersymmetric vacua in which part of the gauge group is unbroken. They can be described in the brane picture and in the low energy field theory in a similar way.

In section 2 we discussed a set of vacua which is described in the brane picture by the curved fivebrane (6), (12) in the near-horizon geometry of N infinite $D4$-branes. We noted that the shape of the curved fivebrane involves in an important way g_s effects. Thus, the low energy field theory description should involve quantum effects in the gauge theory. We next describe these vacua from the gauge theory point of view.

We start with the $U(N+p)$ gauge theory described in the beginning of this subsection. To study its quantum dynamics, it is convenient to add to the theory two $N \times N$ massive gauge singlet matrix chiral superfields M_{LR}, M_{RL}, and replace (27) by

$$W_\lambda = -\frac{1}{\lambda} M_{LR} M_{RL} + M_{LR} \tilde{Q}_R \cdot Q_L + M_{RL} \tilde{Q}_L \cdot Q_R \ . \tag{31}$$

At low energies, we can integrate out the massive gauge singlets. Their equations of motion set

$$M_{LR} = \lambda \tilde{Q}_L \cdot Q_R, \qquad M_{RL} = \lambda \tilde{Q}_R \cdot Q_L; \tag{32}$$

plugging this in (31) leads to (27). Thus, the theory with superpotential (31) is equivalent to the original one (27) at low energies.

Consider now the theory (31). We are interested in vacua in which the singlets M_{LR}, M_{RL} are non-singular matrices. Thus, the quarks Q, $\tilde{Q}$ are massive, and can be integrated out. This gives an effective superpotential for M_{LR}, M_{RL}, which can be calculated as follows. First, we use the scale matching relation between the scale Λ of the theory with $2N$ flavors Q, $\tilde{Q}$ and the scale Λ_L of the pure gauge theory without them, which takes the following form in standard conventions:

$$\Lambda^{N+3p} \det(M_{LR}) \det(M_{RL}) = \Lambda_L^{3(N+p)} \ . \tag{33}$$

We then use the non-perturbative superpotential of the low energy pure gauge theory,

$$W = (N+p)\Lambda_L^3 \ . \tag{34}$$

Combining this with the classical term in (31) we find the full superpotential for M_{LR}, M_{RL}:

$$W = -\frac{1}{\lambda} M_{LR} M_{RL} + (N+p) \left(\Lambda^{N+3p} \det(M_{LR}) \det(M_{RL}) \right)^{\frac{1}{N+p}} \ . \tag{35}$$

Varying with respect to (say) M_{LR}, and using the fact that the matrices M_{LR}, M_{RL} are non-degenerate, we find that

$$M_{LR} M_{RL} = \lambda \left(\Lambda^{N+3p} \det(M_{LR}) \det(M_{RL}) \right)^{\frac{1}{N+p}} \ . \tag{36}$$

Solving this for the determinants, we find

$$\det(M_{LR}) \det(M_{RL}) = \lambda^{\frac{N(N+p)}{p}} \Lambda^{\frac{N(N+3p)}{p}} \ . \tag{37}$$

In the vacua (36), the $SU(N)_L \times SU(N)_R$ global symmetry is spontaneously broken to the diagonal $SU(N)$ by the expectation value of the mesons M_{LR}, M_{RL}. This gives rise to a moduli space of vacua labeled by the expectation value of the associated Nambu-Goldstone bosons (which are linear combinations of the fields in M_{LR} and M_{RL}).

We found that the quantum effects lead in this case to two (related) phenomena. One is that the chiral superfields $Q_{L,R}$, $\tilde{Q}_{L,R}$ become massive; their mass matrix, (M_{LR}, M_{RL}), is given by (36), (37). The other is the non-zero expectation value of $\tilde{Q}_L \cdot Q_R$, which breaks the gauge symmetry $U(N+p) \to U(p)$. The scale of the breaking is proportional to the gauge coupling and to the expectation value of Q, $\tilde{Q}$. Depending on the parameters of the theory (Λ, λ, g_{YM}) one of the effects occurs at a higher energy and dominates the dynamics.

If the scale of the breaking of the gauge symmetry is high, the low energy theory can be thought of as a pure $U(p)$ $\mathcal{N}=1$ gauge theory with massive flavors. The scale of this theory can be obtained by plugging (36) into (35), which gives

$$\Lambda_p^3 = \left(\Lambda^{N+3p}\lambda^N\right)^{\frac{1}{p}} \ . \tag{38}$$

In the brane system, the role of Λ_p is played by ξ/l_s^2 (6).

The spectrum of the above gauge theory contains mesons whose masses are difficult to compute, since this is non-holomorphic information. In section 4 we will see that in the regime where the brane description is valid, the masses of the flavor-singlet mesons can be computed using the gravitational description.

3.2 Compact x_6

As mentioned above, another interesting brane configuration is obtained by compactifying the x_6 direction on a circle, as in figure 5. We discuss here only the supersymmetric case of periodic boundary conditions for the fermions on the circle. The low-energy gauge theory in this case is an $SU(N+p)\times SU(N)$ four dimensional $\mathcal{N}=1$ supersymmetric gauge theory, with two chiral multiplets A_i ($i=1,2$) in the bi-fundamental representation and two (B_i, $i=1,2$) in the anti-bi-fundamental [15, 16, 17, 6]. There is also a superpotential generalizing (27), of the form

$$W = \lambda \mathrm{tr}(A_1B_1A_2B_2 - A_1B_2A_2B_1). \tag{39}$$

One can discuss this brane configuration in a decoupling limit which preserves the five dimensional gauge dynamics including the Kaluza-Klein modes on the circle, or take a different decoupling limit that keeps only the four dimensional gauge dynamics. The latter limit was discussed in [6], but we expect the moduli space to be independent of precisely which limit we take. As argued in [6] (see [18] for a review), as one decreases the energy, this theory undergoes a series of "duality cascades", such that the effective field theory describing physics at lower energy scales is first a $SU(N)\times SU(N-p)$ theory, then a $SU(N-p)\times SU(N-2p)$ theory, and so on. In the gravitational solution studied in [6], this cascade eventually ends (if N is a multiple of p) in a confining background. This type of renormalization group flow implies that also at high energies we cannot (once we take a decoupling limit from string theory) discuss the theory at fixed N, since this theory has a Landau pole, but we have to increase N as we increase the UV cutoff, and if we take the UV cutoff to infinity we must also take N to infinity at the same time.

The moduli space of this theory was analyzed in detail in [11], and we will just quote their results here. They found that this theory actually has a large moduli space of vacua. When N is not a multiple of p (and $p \ll N$), but rather $N \bmod p = q$, the theory has branches of the moduli space of (real) dimensions $6q$, $6(q+p)$, $6(q+2p)$, $\cdots$, $6(q+np)$, $\cdots$, which are each equivalent to a symmetric product of deformed conifolds (when the UV cutoff goes to infinity, there is an infinite number of branches of this type). When $q = 0$, the only difference is that the dimension of the lowest branch is not zero but two, and on this lowest branch some baryonic operators condense.

Seeing this moduli space in the Klebanov-Strassler description, corresponding to the decoupling limit of the four dimensional gauge theory when this theory is strongly coupled, is straightforward but not completely trivial; the different branches involve $(q+np)$ $D3$-branes that are free to move on the conifold and that generally back-react on its geometry. We can see precisely the same moduli space in the gravitational description of the previous section, which is valid in the opposite limit in which the five dimensional gauge dynamics is strongly coupled. We saw in the previous section that in this description the background involves a spiralling fivebrane, which reduces the rank of the gauge group as we go down in the radial direction, just as in the Klebanov-Strassler cascade (but here this reduction comes from explicit brane sources, rather than from background fluxes). We also saw that for given UV boundary conditions we have an infinite set of vacua (24), which differ in the value of the extreme IR flux (we identify the vacuum in which the IR flux is equal to $(q+np)$ with the $6(q+np)$-dimensional branch of the moduli space mentioned above). Note that, unlike in the case of Klebanov and Strassler, in our case we do not have a good description of the vacua with the lowest values of n, where back-reaction of the five-brane becomes important in our solutions. However, we do have good descriptions for all the vacua with $p \ll q + np$.

4 Observables and spectra

In the decoupling limit described in section 2.2, holography relates non-normalizable modes in the bulk to operators in the dual field theory, and the bulk path integral with sources to the generating functional of correlation functions of the corresponding field theory operators. Normalizable states in the bulk correspond to states in the dual field theory.

In this section we discuss some aspects of this map for operators and states associated with the brane intersection, which correspond to fields living on the curved $NS5$-brane.[12] We start by identifying modes of the fivebrane with operators in the low energy field theory, and then move on to calculate the spectrum of mesons. In section 4.4 we discuss the potential existence of normalizable Nambu-Goldstone modes associated with global symmetry breaking.

---

[12] There are also "bulk" modes associated with the field theory living on the infinite $D4$-branes. In the probe approximation (*i.e.* to leading order in p/N) they are not affected by the extra fivebrane.

4.1 Operator matching

The curved $NS5$-brane (6) has a two component boundary, corresponding to large positive x_6 (and hence large w (8)), and large negative x_6 (and large v). The latter corresponds to going to the boundary at large u along the NS-brane; the former, along the NS'-brane. We will focus, for concreteness, on the operator map for modes defined on the NS-brane. The discussion of modes living on the NS'-brane is very similar.

To get a qualitative guide to the spectrum of operators on the NS-brane, it is convenient to go back to figure 3, which does not take into account g_s effects, but is nevertheless useful. In this limit, the NS-brane fills the v plane; the bosonic modes living on it are scalar fields describing its fluctuations in the transverse directions[13] (w, z, x_7), and a self-dual 2-form field. Translations of the fivebrane in z change the coupling of the $U(N+p)$ gauge theory. Hence, the bulk operator corresponding to the relative location of the NS and NS'-branes couples to the gauge theory Lagrangian. We are mainly interested in mesons, and will not discuss this mode in detail.

The scalar fields w and x_7 couple to gauge-invariant operators constructed out of the chiral superfields Q_L, $\tilde{Q}_L$ living at the intersection of the $D4$-branes and the NS-brane. To study these operators in the bulk, we write

$$x_7 = \epsilon \tilde{x}_7(v, \bar{v}, x^\mu), \quad w = w_0(v) + \epsilon \tilde{w}(v, \bar{v}, x^\mu), \tag{40}$$

where w_0 is the undeformed profile (6), and we work to leading order in the deformations $\tilde{x}_7$, $\tilde{w}$. Specializing to wavefunctions with well defined charge (9),

$$\tilde{w} = v^n \tilde{w}^{(n)}(v\bar{v}, x^\mu), \quad \tilde{x}_7 = v^n \tilde{x}_7^{(n)}(v\bar{v}, x^\mu), \tag{41}$$

and plugging into the fivebrane action, we find the linearized equation of motion

$$\left(\partial_{x_6}^2 + \partial_\mu \partial^\mu\right) F - \frac{n^2}{\lambda_p^2} F = 0, \tag{42}$$

where $F = \tilde{w}^{(n)}, \tilde{x}_7^{(n)}$, and we used x_6 rather than v to parameterize the worldvolume of the unperturbed fivebrane, (6).

The field theory operators corresponding to (41) can be identified by matching the transformation properties under the symmetries[14]. This leads to

$$\begin{aligned} v^n \tilde{w}^{(n)} &\leftrightarrow Q_L \Phi_L^n \tilde{Q}_L, \\ v^n \tilde{x}_7^{(n)} &\leftrightarrow Q_L \Phi_L^n Q_L^\dagger - \tilde{Q}_L^\dagger \Phi_L^n \tilde{Q}_L. \end{aligned} \tag{43}$$

---

[13] From the ten dimensional point of view, one of these scalar fields, $\mathrm{Im}(z) = x_{11}$, is non-geometric.

[14] Of course, all the states and operators we discuss are singlets of the $SU(N)_L \times SU(N)_R$ symmetry, since this is a gauge symmetry in our setup.

So far, we have focused on operators associated with the boundary along the NS-brane, at large negative x_6. Of course, perturbations introduced there propagate to large positive x_6 (where the equations of motion (42) are corrected in a way that will be described below). In order to study perturbations containing only the operators (43), and not their analogs with $L \to R$, one needs to choose solutions of the equations of motion that decay rapidly as $x_6 \to \infty$.

At first sight, for $n > 0$ the operators on the right-hand side of (43) seem to involve the five dimensional modes living on the infinite $D4$-branes. However, one can use the equations of motion of Q, $\tilde{Q}$ (29) to express them purely in terms of fields in the four dimensional low energy theory. For example, the $\tilde{w}^{(1)}$ mode corresponds to the operator $\tilde{Q}_L \cdot Q_R \tilde{Q}_R \cdot Q_L$. For higher n, one finds operators of the schematic form $\tilde{Q}_L Q_L (\tilde{Q}_R Q_R)^n$. A similar set of operators with $L \leftrightarrow R$ is obtained from the other boundary, $x_6 \to \infty$.

To analyze the solutions of (42), it is convenient to write the operators (43) in momentum space, *i.e.* take $F(x_6, x_\mu) = F(x_6) \exp(ik^\mu x_\mu)$. For $n > 0$ and sufficiently small k^2, the solutions of (42) are in general non-normalizable as $x_6 \to -\infty$; this gives rise to the sources holographically related to the operators in the low energy theory at the intersection, (43). For $n = 0$ and timelike (or null) momentum, $k^2 \leq 0$, the solutions of (43) are in fact delta-function normalizable; they correspond to $4 + 1$ dimensional scattering states which are not localized at the intersection. The same is true for non-zero n and $-k^2 \geq n^2/\lambda_p^2$. Thus, we conclude that the map (43) is only valid for $n \geq 1$, and that in addition to the field theory modes, the brane system contains a continuum of four dimensional states above the gap n/λ_p. For $n = 0$, this continuum starts at zero energy.[15]

Having a continuum with some discrete localized states is natural in field theories with defects. In our case we can think of the brane intersection region as a codimension one defect inside the fivebrane; the discrete states are localized near the intersection, while the continuum corresponds to states propagating in the bulk of the fivebrane. In the limit we take, the four dimensional fields are not decoupled from the higher dimensional ones.

The continuous spectrum in our system is also somewhat similar to what happens in Little String Theory, where in a given charge sector one typically finds a discrete spectrum of localized modes and a continuous spectrum corresponding to modes propagating in an asymptotically linear dilaton throat (see *e.g.* [19]). However, in that case the continuum does not have a simple interpretation in terms of a local field theory on the fivebranes.

In the next subsection we turn to the spectrum of normalizable modes of the confining vacuum (6) of holographic MQCD. These modes correspond to particles in the dual gauge theory. We will focus on the scalars corresponding to transverse fluctuations of the fivebrane. Parametrizing the worldvolume as in section 2, we can label these directions by v, w and x_7. Since the classical shape (6) is localized at $x_7 = 0$, it is easiest to study the fluctuations in the x_7 direction, which are decoupled from those in the other directions. Thus, we start with those, and then move on to the other ones.

---

[15] The fact that the map (43) is not valid for $n = 0$ is natural from the field theory perspective, since the operators on the right-hand side of (43) are analogs of the $U(1)$ part of $U(N)$ in other holographic systems.

We find a discrete spectrum of massive states, below the continuum discussed above. If we take a limit that decouples the higher dimensional fields from the four dimensional field theory, the continuous part of the spectrum should decouple, leaving behind a set of discrete states. We assume that the discrete states in this limit are continuously related to the ones we find here, but it is also possible that as we interpolate between the two regimes, some states disappear into the continuum while other states emerge from it.

4.2 Spectrum of x_7 fluctuations

We start with the analysis of the fluctuations along the x_7 direction which is transverse to the $D4$, NS and NS'-branes. As mentioned above, these fluctuations are decoupled from those corresponding to the other directions, and therefore their analysis is more tractable. The corresponding gauge theory meson operators are described in the previous subsection (see (43) and the discussion around it).

Before incorporating the fluctuations, the x_7 position of the fivebrane is a constant, $x_7 = X_7$ (which is one of the components of $\vec{r_0}$ in (14)). Perturbing around the classical solution, we have

$$x_7 = X_7 + \epsilon \tilde{x}_7(z, \bar{z}, x^\mu). \tag{44}$$

Consider first the case $X_7 = 0$. Plugging (44) into the background (14)

$$\begin{aligned} ds^2 &= H^{-1/3}[dx_\mu^2 + dz d\bar{z}] + H^{2/3}[dv d\bar{v} + dw d\bar{w} + dx_7^2], \\ C_6 &= \frac{1}{2i} H^{-1} d^4x \wedge dz \wedge d\bar{z}, \qquad H = \frac{\pi \lambda_N l_s^2}{(v\bar{v} + w\bar{w} + x_7^2)^{3/2}}\,, \end{aligned} \tag{45}$$

we find the induced metric on the probe branes,

$$ds^2 = H^{-1/3}\left(dx_i^2 + g_{ab} dx^a dx^b\right), \qquad x^a = (z, \bar{z}, t). \tag{46}$$

In (46) and below we take (without loss of generality) the field $\tilde{x}_7$ to depend only on the time t and not on the spatial coordinates in $\mathbb{R}^{3,1}$. Keeping only terms up to quadratic order in ϵ, we find (denoting $\partial \equiv \partial/\partial z$, $\bar{\partial} = \partial/\partial \bar{z}$)

$$g = \begin{pmatrix} \epsilon^2 H \partial\tilde{x}_7 \partial\tilde{x}_7 & g_{z\bar{z}} & \epsilon^2 H \partial\tilde{x}_7 \dot{\tilde{x}}_7 \\ g_{z\bar{z}} & \epsilon^2 H \bar{\partial}\tilde{x}_7 \bar{\partial}\tilde{x}_7 & \epsilon^2 H \bar{\partial}\tilde{x}_7 \dot{\tilde{x}}_7 \\ \epsilon^2 H \partial\tilde{x}_7 \dot{\tilde{x}}_7 & \epsilon^2 H \bar{\partial}\tilde{x}_7 \dot{\tilde{x}}_7 & -1 + H\epsilon^2 (\dot{\tilde{x}}_7)^2 \end{pmatrix}. \tag{47}$$

Here we defined

$$g_{z\bar{z}} = \frac{1}{2}[1 + H(|\partial v|^2 + |\partial w|^2)] + \epsilon^2 H |\partial \tilde{x}_7|^2.$$

The harmonic function takes the form

$$H = \frac{\pi \lambda_N l_s^2}{(u_{cl}^2 + x_7^2)^{3/2}} = \frac{\pi \lambda_N l_s^2}{\left[2\xi^2 \cosh\left(\frac{2x_6}{\lambda_p}\right) + x_7^2\right]^{3/2}} . \tag{48}$$

We also define the dimensionless coordinates $\hat{x}_7 = \frac{\tilde{x}_7}{\sqrt{2}\xi}$, $\hat{x}_6 = \frac{x_6}{\lambda_p}$, $\hat{x}_{11} = \frac{x_{11}}{\lambda_p}$, $\hat{t} = \frac{t}{\lambda_p}$, and the dimensionless ratio q

$$q \equiv \frac{\sqrt{2}\xi \lambda_p^2}{\pi \lambda_N l_s^2} . \tag{49}$$

Note that the condition (22) implies that $q \gg (\lambda_p/\lambda_N)^2 = (p/N)^2$. However, this lower bound on q is very weak in the probe limit $p \ll N$. E.g., if we keep the "QCD scale" ξ/l_s^2 fixed in units of the five dimensional gauge coupling λ_p or λ_N, we have $q \ll 1$ in the probe limit.

In terms of the above definitions and omitting the hats, expanding to second order in ϵ we find in this case the following Lagrangian (up to a multiplicative constant)

$$\begin{aligned} L &\simeq \int d^2 z H^{-1} \left(\sqrt{\frac{1}{2}\det(2g)} - 1\right) \simeq \\ &\simeq \epsilon^2 \int d^2 z \left[2\partial x_7 \bar{\partial} x_7 - \frac{1}{2}\left(1 + \frac{1}{q\sqrt{\cosh(2x_6)}}\right) \dot{x}_7^2 \right] . \end{aligned} \tag{50}$$

The corresponding equation of motion for $x_7(x_6, x_{11}, t)$ is

$$-\partial_{x_6}^2 x_7 - \partial_{x_{11}}^2 x_7 + \left(1 + \frac{1}{q\sqrt{\cosh(2x_6)}}\right) \ddot{x}_7 = 0. \tag{51}$$

If we look at a mode of fixed mass and momentum in x_{11}, such that $x_7(x_6, x_{11}, t) = e^{inx_{11} + imt} x_7(x_6)$, we get the following equation[16]

$$\partial_{x_6}^2 x_7 - \left[n^2 - m^2 \left(1 + \frac{1}{q\sqrt{\cosh(2x_6)}}\right)\right] x_7 = 0. \tag{52}$$

In the asymptotic region of large $|x_6|$, the equation we find is (42)

$$\partial_{x_6}^2 x_7 = (n^2 - m^2) x_7 . \tag{53}$$

Clearly, for $m^2 > n^2$ the solutions in this region are just plane waves, while for $m^2 < n^2$ the solutions decay or grow exponentially at infinity. We can think of (52) as a

---

[16] Note that the charge of the state under the global $U(1)$ symmetry (9) is $-n$.



Figure 8: The spectrum of x_7 fluctuations: m^2 (in units of $1/\lambda_p^2$) as a function of q.

Schrödinger equation describing the motion of a particle with vanishing energy in the potential

$$V(x_6) = n^2 - m^2 \left(1 + \frac{1}{q\sqrt{\cosh(2x_6)}}\right). \tag{54}$$

This makes it clear that when $m^2 > n^2$ the solutions describe scattering off a potential well and there is a delta-function normalizable solution for any value of m^2, while for $m^2 < n^2$ the solutions describe bound states in a potential so we expect to get solutions only for discrete values of m^2 (one can show that at least one bound state exists for all q and n).

Going back to dimensionful variables, we see that we have a continuum of states with $m > n/\lambda_p$, and a discrete spectrum for $m < n/\lambda_p$. As discussed above, the discrete spectrum can be thought of as describing mesons in the confining vacuum of the gauge theory of section 3 at strong coupling, while the continuum is associated with higher dimensional modes.

To examine the spectrum of mesons, we numerically solved (51) for the case $n = 1$. The resulting spectrum is depicted in figure 8 as a function of q. The qualitative features of this spectrum are obvious by thinking about the analogous Schrödinger problem. For small q, the potential well (54) is very deep, so we expect many bound states, with the low lying ones having $m^2 \propto q/\lambda_p^2$. On the other hand, for large q the potential is very shallow, so we expect (and find) just a single bound state, very close to $m = 1/\lambda_p$.

Note that the potential (54) is symmetric under $x_6 \to -x_6$, which acts as charge conjugation on the vector superfields, and exchanges $Q_L \leftrightarrow Q_R$. The minimum of the potential is at $x_6 = 0$, and it monotonically increases with $|x_6|$. The mesons depicted in figure 8 are localized near $x_6 = 0$; they can be thought of as having significant overlap with both Q_L and Q_R.

For $n > 1$ the states described by (52) can be thought of as "exotics" since they have the $U(1)$ charge of $Q_L^\dagger Q_L (\tilde{Q}_R Q_R)^n$. Their spectrum is similar to that of figure 8, but the masses are larger. This can be seen by noting that to have zero energy bound states, the potential (54) has to be negative at the origin. This implies that the bound state masses m always satisfy the bound

$$(m\lambda_p)^2 > \frac{n^2}{1+\frac{1}{q}} \; . \tag{55}$$

Thus, the masses of "exotics" containing n pairs of $\tilde{Q}_R Q_R$ and/or $\tilde{Q}_L^\dagger Q_L^\dagger$ grow with n.

For the "massive" case, where the N $D4$-branes are displaced from the p $D4$-branes that stretch between the NS and NS'-branes in the x_7 direction by the distance X_7 (see figure 2), the harmonic function takes the form

$$H = \frac{\pi \lambda_N l_s^2}{\left[2\xi^2 \cosh\left(\frac{2x_6}{\lambda_p}\right) + (X_7 + \epsilon \tilde{x}_7)^2\right]^{3/2}} \; . \tag{56}$$

The equation of motion reads (after rescaling as above, with X_7 rescaled in the same way as $\tilde{x}_7$)

$$\partial_{x_6}^2 x_7 - \left[n^2 - m^2\left(1 + \frac{\cosh(2x_6)}{q(\cosh(2x_6) + X_7^2)^{3/2}}\right)\right] x_7 = 0. \tag{57}$$

The potential (54) takes in this case the form

$$V(x^6) = n^2 - m^2\left(1 + \frac{\cosh(2x_6)}{q(\cosh(2x_6) + X_7^2)^{3/2}}\right). \tag{58}$$

For $X_7 < 1/\sqrt{2}$ or, in terms of dimensionful variables, $X_7 < \xi$, the potential (58) is qualitatively similar to (54) – it has a unique minimum at $x_6 = 0$ and no other extrema. For $X_7 > 1/\sqrt{2}$, the extremum at $x_6 = 0$ becomes a local maximum, and the potential (58) becomes a double well potential, with minima at the two solutions of the equation $\cosh(2x_6) = 2X_7^2$. As X_7 increases, the potential becomes more and more sharply peaked (for sufficiently small q), and the two minima move to large $|x_6|$.

For large X_7 and small q, the solutions of the Schrödinger problem that gives the masses, (57), split into wavefunctions localized in the two wells. If this was the end of the story, the theory would break the charge conjugation symmetry $x_6 \to -x_6$, and the mass spectrum would split into degenerate doublets related by the symmetry.

Of course, as is well known, symmetry breaking does not happen in quantum mechanics due to tunneling. Rather than exactly degenerate doublets, we expect to find approximately degenerate pairs of states corresponding to the sum and difference of wavefunctions localized in the left and right wells. In figures 9-11 we present numerical results for the mass spectrum for three values of X_7 in the "broken phase", $X_7^2 = 1, 10, 100$. As one can see from these figures, for fixed X_7 the splitting of the spectrum into approximately degenerate pairs becomes more and more pronounced as q decreases, while for fixed q it becomes more pronounced as X_7 increases, as one would expect.



Figure 9: The spectrum of x_7 fluctuations for $X_7^2 = 1$.



Figure 10: The spectrum of x_7 fluctuations for $X_7^2 = 10$.



Figure 11: The spectrum of x_7 fluctuations for $X_7^2 = 100$.

From the point of view of the low energy field theory of section 3, the change in the nature of the wavefunctions of the normalizable states described above means that as X_7 increases, the mesons go from having significant overlaps with both Q_L and Q_R to being primarily made out of Q_L (for wavefunctions localized in the left well) and of Q_R (for those in the right well). These "flavor eigenstates" have small mixings, such that the mass eigenstates are the sum and difference of the mesons made out of Q_L and those made out of Q_R.

4.3 Fluctuations of the other transverse directions

Next, we analyze the transverse fluctuations in some of the other directions. From the field theory point of view, one expects to find a qualitatively similar spectrum to that described in the previous subsection. As we will see, in this case the equations for the fluctuations are coupled, and thus are more difficult to solve numerically. However, in certain limits we will be able to solve them, and find that the expectations are realized.

As before, to study the small fluctuations we expand around the holomorphic profile,

$$w = w_0(z) + \epsilon w_1(z, \bar{z}, t), \qquad v = v_0(z) + \epsilon v_1(z, \bar{z}, t). \tag{59}$$

Similar to (46) we determine the nontrivial part of g_{ab} in the induced metric. To compute the determinant of g_{ab} to quadratic order in ϵ, we should keep quadratic terms in $g_{z\bar{z}}$ and g_{tt}, and linear contributions to the other components of g_{ab}:

$$g = \begin{pmatrix} \epsilon H(\partial w_0 \partial \bar{w}_1 + \partial v_0 \partial \bar{v}_1) & g_{z\bar{z}} & \frac{H\epsilon}{2}(\partial w_0 \dot{\bar{w}}_1 + \partial v_0 \dot{\bar{v}}_1) \\ g_{z\bar{z}} & \epsilon H(\bar{\partial} \bar{w}_0 \bar{\partial} w_1 + \bar{\partial} \bar{v}_0 \bar{\partial} v_1) & \frac{H\epsilon}{2}(\bar{\partial} \bar{w}_0 \dot{w}_1 + \bar{\partial} \bar{v}_0 \dot{v}_1) \\ \frac{H\epsilon}{2}(\partial w_0 \dot{\bar{w}}_1 + \partial v_0 \dot{\bar{v}}_1) & \frac{H\epsilon}{2}(\bar{\partial} \bar{w}_0 \dot{w}_1 + \bar{\partial} \bar{v}_0 \dot{v}_1) & H\epsilon^2(\dot{w}_1 \dot{\bar{w}}_1 + \dot{v}_1 \dot{\bar{v}}_1) - 1 \end{pmatrix}.$$

Here

$$2g_{z\bar{z}} = 1 + H[\partial(w_0 + \epsilon w_1)\bar{\partial}(\bar{w}_0 + \epsilon \bar{w}_1) + \partial(v_0 + \epsilon v_1)\bar{\partial}(\bar{v}_0 + \epsilon \bar{v}_1) + \epsilon^2 \bar{\partial} w_1 \partial \bar{w}_1 + \epsilon^2 \bar{\partial} v_1 \partial \bar{v}_1]. \tag{60}$$

Next we compute

$$\begin{aligned} \det(2g) &= 2\Big[(2g_{z\bar{z}})^2(-g_{tt}) - 4(\epsilon H)^2(\bar{\partial}\bar{w}_0 \bar{\partial} w_1 + \bar{\partial}\bar{v}_0 \bar{\partial} v_1)(\partial w_0 \partial \bar{w}_1 + \partial v_0 \partial \bar{v}_1)\Big] \\ &\quad + 2(\epsilon H)^2(\partial w_0 \dot{\bar{w}}_1 + \partial v_0 \dot{\bar{v}}_1)(\bar{\partial}\bar{w}_0 \dot{w}_1 + \bar{\partial}\bar{v}_0 \dot{v}_1)(2g_{z\bar{z}}), \end{aligned} \tag{61}$$

so that

$$\begin{aligned} \sqrt{\frac{1}{2}\det(2g)} &= (2g_{z\bar{z}})[1 - \frac{H\epsilon^2}{2}(\dot{w}_1 \dot{\bar{w}}_1 + \dot{v}_1 \dot{\bar{v}}_1)] \\ &\quad + \frac{2(\epsilon H)^2}{1 + H(\partial w \bar{\partial} \bar{w} + \partial v \bar{\partial} \bar{v})}\Big[-(\bar{\partial}\bar{w}_0 \bar{\partial} w_1 + \bar{\partial}\bar{v}_0 \bar{\partial} v_1)(\partial w_0 \partial \bar{w}_1 + \partial v_0 \partial \bar{v}_1)\Big] \\ &\quad + \frac{(\epsilon H)^2}{2}(\partial w_0 \dot{\bar{w}}_1 + \partial v_0 \dot{\bar{v}}_1)(\bar{\partial}\bar{w}_0 \dot{w}_1 + \bar{\partial}\bar{v}_0 \dot{v}_1). \end{aligned} \tag{62}$$

The Lagrangian density in $\mathbb{R}^{3,1}$ becomes

$$\begin{aligned} L &= \int d^2z \frac{1}{H}(\sqrt{\frac{1}{2}\det(2g)} - 1) \\ &= \int d^2z \Big[\partial(w_0 + \epsilon w_1)\bar\partial(\bar w_0 + \epsilon \bar w_1) + \partial(v_0 + \epsilon v_1)\bar\partial(\bar v_0 + \epsilon \bar v_1) - \epsilon^2 g_{z\bar z}(\dot w_1 \dot{\bar w}_1 + \dot v_1 \dot{\bar v}_1) \\ &\qquad - \frac{2\epsilon^2 H}{1 + H(\partial w \bar\partial \bar w + \partial v \bar\partial \bar v)}(\bar\partial \bar w_0 \bar\partial w_1 + \bar\partial \bar v_0 \bar\partial v_1)(\partial w_0 \partial \bar w_1 + \partial v_0 \partial \bar v_1) \\ &\qquad + \frac{\epsilon^2 H}{2}(\partial w_0 \dot{\bar w}_1 + \partial v_0 \dot{\bar v}_1)(\bar\partial \bar w_0 \dot w_1 + \bar\partial \bar v_0 \dot v_1) + \epsilon^2 \bar\partial w_1 \partial \bar w_1 + \epsilon^2 \bar\partial v_1 \partial \bar v_1\Big]. \end{aligned} \tag{63}$$

The terms linear in ϵ do not contribute to the equations of motion for w_1 and v_1, and since H appears only multiplying ϵ^2, we can replace it by its value H_0 in the original solution. Dropping an overall power of ϵ, we find

$$\begin{aligned} L &= \int d^2z \Big[\partial w_1 \bar\partial \bar w_1 + \partial v_1 \bar\partial \bar v_1 + \bar\partial w_1 \partial \bar w_1 + \bar\partial v_1 \partial \bar v_1 - \\ &\qquad \frac{2H_0}{F}(\bar\partial \bar w_0 \bar\partial w_1 + \bar\partial \bar v_0 \bar\partial v_1)(\partial w_0 \partial \bar w_1 + \partial v_0 \partial \bar v_1)\Big] \\ &\qquad -\frac{1}{2}\int d^2z \Big[F(|\dot w_1|^2 + |\dot v_1|^2) - H_0(\partial w_0 \dot{\bar w}_1 + \partial v_0 \dot{\bar v}_1)(\bar\partial \bar w_0 \dot w_1 + \bar\partial \bar v_0 \dot v_1)\Big], \end{aligned} \tag{64}$$

where we defined

$$F \equiv 1 + H_0(\partial w_0 \bar\partial \bar w_0 + \partial v_0 \bar\partial \bar v_0). \tag{65}$$

For the profile (6) we have

$$v_0 = \xi e^{-z/\lambda_p}, \quad w_0 = \xi e^{z/\lambda_p}, \quad H_0 = \frac{\pi \lambda_N l_s^2}{(v_0 \bar v_0 + w_0 \bar w_0)^{3/2}}, \quad F = 1 + \frac{2\xi^2}{\lambda_p^2} H_0 \cosh\left(\frac{z + \bar z}{\lambda_p}\right). \tag{66}$$

The equations of motion that follow from (64) are complicated. To get some insight about the structure of the spectrum we first analyze the asymptotic form of the equations of motion at large values of $|\mathrm{Re}(z)| = |x_6|$. Keeping only the leading order terms, we arrive at an approximate asymptotic Lagrangian:

$$L \approx \int d^2z \left[\partial w_1 \bar\partial \bar w_1 + \partial v_1 \bar\partial \bar v_1 + \bar\partial w_1 \partial \bar w_1 + \bar\partial v_1 \partial \bar v_1\right] - \frac{1}{2}\int d^2z \left[|\partial_\mu w_1|^2 + |\partial_\mu v_1|^2\right], \tag{67}$$

where for completeness we have restored the full x_μ dependence. The corresponding equations of motion read

$$\partial\bar\partial w_1 - \frac{1}{4}\partial^\mu \partial_\mu w_1 = 0, \qquad \partial\bar\partial v_1 - \frac{1}{4}\partial^\mu \partial_\mu v_1 = 0. \tag{68}$$

Taking a fixed momentum n in x_{11} and a fixed four dimensional mass m as above, the resulting equation of motion is

$$\partial_{x_6}^2 w_1 - \left(\frac{n^2}{\lambda_p^2} - m^2\right) w_1 = 0, \tag{69}$$

and similarly for the other modes. Thus, as in the previous section, we have a discrete spectrum for $m < |n|/\lambda_p$, and a continuum for $m > |n|/\lambda_p$. Note that since in this case v and w themselves carry a $U(1)$ charge ± 1, these fluctuations carry $U(1)$ charges $n \pm 1$.

Let us describe in more detail the modes coming from fluctuations of the absolute values of v and w, namely

$$V = |v| = V_0 + \epsilon V_1, \qquad W = |w| = W_0 + \epsilon W_1. \tag{70}$$

In terms of the coordinates u and α defined in (11)

$$u e^{i\alpha} = (V_0 + \epsilon V_1) + i(W_0 + \epsilon W_1). \tag{71}$$

Substituting these variables into the Lagrangian (64) we get

$$\begin{aligned} L &= \frac{1}{2}\int dx_6 \left[(V_1')^2 + (W_1')^2 + (V_1^2 + W_1^2) - H_0 \frac{[V_0' V_1' + W_0' W_1' - V_0 V_1 - W_0 W_1]^2}{(\lambda_p^2 + u^2 H)_0} \right] \\ &\quad + \int dx_6 \frac{1}{2}(\lambda_p^2 + u^2 H)_0 [(\partial_\mu V_1)^2 + (\partial_\mu W_1)^2]. \end{aligned} \tag{72}$$

Here we have (after rescaling x_6 as before)

$$V_0 = \xi e^{-x_6}, \quad W_0 = \xi e^{x_6}, \quad (\lambda_p^2 + u^2 H)_0 = \lambda_p^2 + \frac{Q}{u} = \lambda_p^2 + \frac{Q}{\xi (e^{2x_6} + e^{-2x_6})^{1/2}}.$$

Assuming that $W_1 \sim \sin(mt)$, $V_1 \sim \sin(mt)$, we find the Lagrangian

$$\begin{aligned} L &= \frac{1}{2}\int dx_6 \left[(V_1')^2 + (W_1')^2 + (V_1^2 + W_1^2) - [\lambda_p^2 + u^2 H]_0 [(mV_1)^2 + (mW_1)^2] \right] \\ &\quad - \frac{1}{2}\int dx_6 \frac{H_0}{(\lambda_p^2 + u^2 H)_0} [W_0 W_1' - V_0 V_1' - V_0 V_1 - W_0 W_1]^2. \end{aligned} \tag{73}$$

This Lagrangian is still too complicated to analyze. In the Appendix we invoke several approximations that enable us to simplify its form for some range of parameters. Using these approximations we get a decoupled Lagrangian for $V_+ \equiv \frac{1}{\sqrt{2}}(V_1 + W_1)$ of the form

$$L \approx \frac{1}{2}\int dx_6 \left[(V_+')^2 + \left(1 - (m\lambda_p)^2 - \frac{(m\lambda_p)^2}{q} \frac{1}{\sqrt{\cosh(2x_6)}} \right) V_+^2 \right]. \tag{74}$$

This is exactly the same Lagrangian as the one we found (with no assumptions and approximations) for the fluctuations along x_7 (50). The solution of the corresponding equation of motion is thus identical (for $n = 1$) to the numerical solution described in figure 8.

4.4 Self-dual B field

The remaining bosonic field living on the curved fivebrane is the self-dual 2-form B field. Its fluctuations give rise to a spectrum of mesons, whose masses can be analyzed as in the previous subsections. We will not discuss the details of this analysis here.

Instead, we will comment briefly on the following question. In the original brane configuration of figure 3, there are in fact two independent self-dual B fields, living on the two $NS5$-branes. One can think of them as generating two $U(1)$ symmetries in the low energy field theory. The corresponding gauge fields are obtained by reducing the self-dual B-field living on the NS-brane on the angular direction in the v-plane, and similarly for the NS'-brane.

In the confining vacuum described by (6), the two $NS5$-branes connect, and these symmetries are broken to the diagonal subgroup, $U(1)_L \times U(1)_R \to U(1)_D$. Superficially, the situation seems to be very similar to that in the Sakai-Sugimoto model [20] (a closer analog to our situation is the non-compact analog of that model, studied in [21]). In that case, the symmetry breaking was spontaneous, and gave rise to a massless pion, which corresponded to the zero mode of the component of the gauge field on the flavor $D8$-branes along the U-shaped brane. In analogy, one might expect that in our system a massless pion would arise from the self-dual B field with components along the curved fivebrane. However, there are two important differences between our case and the Sakai-Sugimoto model. In our case the brane only approaches the boundary at infinite values of x_6; also, in our case the radius of the x_{11} circle in the brane worldvolume goes to infinity when we approach the boundary. Thus, it is not clear if we really have a spontaneously broken global symmetry as described above. To check this we look for a mode of the fluctuation of the $B_{\mu\nu}$ field that corresponds to a massless mode in the dual field theory, and which is normalizable.

Consider the fluctuation modes of the self–dual two-form field B living on the $M5$-brane. Recall first that the induced metric on the $M5$-brane is (see (14)) given by

$$ds^2_{ind} = H^{-1/3}\left[dx_\mu^2 + (1 + H(|\partial v|^2 + |\partial w|^2))dzd\bar{z}\right] . \tag{75}$$

Classically, the self–dual B-field living on the $M5$-brane vanishes. To study a candidate for a Nambu–Goldstone mode, we only excite the components of the B-field which give rise to scalars in spacetime:

$$B = ib\ dzd\bar{z} + (\partial_\mu \phi dx^\mu dz + \text{c.c}) + \hat{B}_{\mu\nu} dx^\mu dx^\nu . \tag{76}$$

All fields appearing here are functions of $(x_\mu, z, \bar{z})$. The field ϕ can be removed by a gauge transformation, and then the self–duality condition determines $\hat{B}_{\mu\nu}$ in terms of b. Focusing on the first term in (76), we find

$$\begin{aligned} &B = ib\ dzd\bar{z}, \quad dB = i(\partial_\mu b)dx^\mu dzd\bar{z}, \quad *dB = ig^{z\bar{z}}H^{-1/3}(*_4 d_4 b), \\ &d(*dB) = ig^{z\bar{z}}H^{-1/3}\partial_\mu\partial^\mu b\ d^4x + id_2[(g^{z\bar{z}}H^{-1/3})(*_4 d_4 b)]. \end{aligned} \tag{77}$$

Since the B-field is self–dual, the expression in the second line must vanish. This implies that b is a massless mode, namely, $\partial^\mu\partial_\mu b = 0$. Let us check whether such a mode is normalizable. The second term in the last equation allows us to determine the dependence of b on $(z, \bar{z})$:

$$b = H^{1/3} g_{z\bar{z}} \tilde{b}(x_\mu) = \left[1 + \frac{H}{\lambda_p^2}(v\bar{v} + w\bar{w})\right] \tilde{b}(x_\mu). \tag{78}$$

To determine the norm of this expression, we should evaluate

$$\int dz d\bar{z} \sqrt{-g}(-g^{tt})(g^{z\bar{z}})^2 (\partial_\mu b)^2 \sim \int dz d\bar{z} (\partial_\mu \tilde{b})^2, \tag{79}$$

which clearly diverges. The expression (79) indicates that, as we found in the previous subsections, the spectrum of this mode is continuous and there is no normalizable four dimensional mode (but just a continuum corresponding to a massless field in six dimensions). Thus, we cannot view the global symmetry discussed above as spontaneously broken, and presumably it is not a good global symmetry of the four dimensional theory for the reasons discussed above. The absence of Nambu-Goldstone bosons was also noted in a similar situation in [22], in a background which also exhibits a continuous spectrum. Note that in the calculation described above we did not explicitly take into account the self–duality constraint. However, we have verified that a careful analysis using the action for a self–dual field presented in [23] leads to the same result.

5 Energy scales

In this section we will discuss the energy scales that enter the dynamics of the brane configurations of section 2. We start with the configuration of figure 1 in weakly coupled type IIA string theory, with ξ, λ_p large in string units. One way to determine the confinement scale in this theory is to calculate the potential between a heavy quark and anti-quark separated in $\mathbb{R}^3$ by a large distance L (not to be confused with the distance in x_6 between the $NS5$-branes, denoted by L in section 2). In QCD, this potential goes at large distances like $V(L) \simeq L T_{\rm conf}$, where $T_{\rm conf}$ is the tension of the QCD string, and the masses of glueballs are of order $\sqrt{T_{\rm conf}}$.

In MQCD, this confining string was discussed in [9]. Although this paper considered the case of large g_s, the construction described in it generalizes trivially to the weakly coupled theory studied here. The confining string can be viewed as an $M2$-brane ending on the $M5$-brane described by (6). At fixed z, this $M5$-brane looks like p points on a circle in the v plane, and p points in the w plane. The locations of the points in the two planes are correlated and change with z. The usual type IIA string is an $M2$-brane wrapped around the x_{11} circle, but if this $M2$-brane ends on the $M5$-brane, it can be continuously deformed to an $M2$-brane which sits at a fixed value of x_{11}, and stretches between two adjacent points in v and w. This string minimizes its energy if it sits at $x_6 = 0$, where its tension is [9] (up to numerical constants)

$$T_{\text{conf}} \simeq \frac{\xi}{p g_s l_s^3} = \frac{\xi}{\lambda_p l_s^2} \,. \tag{80}$$

From the point of view of the brane configuration of figure 1, this confining string (being an $M2$-brane localized in x_{11}) is really a $D2$-brane stretched between the $D4$-branes.

To have a regime in which the dynamics of the theory is dominated by the confining string, the tension (80) must be well below the fundamental type IIA string scale. This is the case if $\xi \ll \lambda_p$. Note that small fluctuations of the fivebrane, of the sort studied in section 4, do not give rise to any normalizable states in flat spacetime, and it is not clear if any discrete states exist.

One may think that when a D-string becomes lighter than the fundamental string, perturbation theory would break down. However, in our case this D-string only exists when it is bound to the fivebrane (which we view as a probe), so it does not affect the validity of perturbation theory in the bulk (or of the probe approximation to the fivebrane physics).

So far, we discussed the system of section 2 for $N = 0$. We now add to the brane configuration $N \gg p$ infinite $D4$-branes, and restrict to their near-horizon geometry. Without the curved fivebrane (6), the quark-anti-quark potential now takes the form $V(L) \propto -\lambda_N/L^2$ [24]. This is obtained by considering the minimal energy type IIA string ending at two points on the boundary separated by a distance L in $\mathbb{R}^3$.

When we add the probe fivebrane (6), the quark-anti-quark potential can in principle change. For very small L, the type IIA string does not reach the curved fivebrane (6) and cannot connect to it, while for very large L the string can go deep into the region of small v, w where its tension goes to zero, and this gives the minimal energy configuration. However, when the minimal radial position of the type IIA string is of order ξ, the confining string could have a smaller energy. Clearly, this can only happen if the tension of the confining string is smaller than the tension of a type IIA string at ξ.

The tension of the confining string (80) is not affected by the $D4$-brane background. However, the fundamental type IIA string tension is corrected. When the N $D4$-branes are at $v = w = 0$, one has (for $x_7 = 0$)

$$H = \frac{g_s N l_s^3}{(|v|^2 + |w|^2)^{\frac{3}{2}}} \,. \tag{81}$$

This harmonic function renormalizes the tension of a type IIA string sitting at some (v, w) to

$$T_{IIA} = \frac{H^{-1/2}}{l_s^2} \,. \tag{82}$$



Figure 12: The quark anti-quark potential $\frac{V(L)\lambda_p^2}{\lambda_N}$ as a function of the distance $\frac{L}{\lambda_p}$. The blue line denotes the potential for a string unattached to the fivebrane, while the purple and green lines are the potentials for attached strings with $q = 0.5, 3$, respectively. For $q = 3$, the actual potential is the minimum between the blue line and the green line, which is drawn in red.

Thus, the ratio between the tensions of the confining string and of a type IIA string at $u \simeq \xi$ is

$$\frac{T_{\text{conf}}}{T_{IIA}} \simeq \left(\frac{\lambda_N l_s^2}{\xi \lambda_p^2}\right)^{\frac{1}{2}} . \tag{83}$$

If

$$\frac{\xi}{l_s^2} \gg \frac{\lambda_N}{\lambda_p^2}, \tag{84}$$

or equivalently $q \gg 1$ in the notations of the previous section, there is a range of L's for which the confining string dominates the quark-antiquark potential. In this range of L's the quark-anti-quark potential is given by $V(L) = T_{\text{conf}} L - c\xi/l_s^2$ for some positive constant c, where the second term comes from the renormalized energy of the type IIA string going down to the $M5$-brane at ξ and then coming back out. The precise quark-anti-quark potential is given (see figure 12) by the minimum between this expression and $V(L) = -\lambda_N/L^2$. For $q \ll 1$ the latter expression is always smaller, but for $q \gg 1$ there is a range of distances where the confining string dominates the quark-anti-quark force.

In section 4 we saw that the spectrum of mesons in holographic MQCD is particularly rich in the opposite limit, $q \ll 1$. The masses of the mesons range in this limit between $\sqrt{q}/\lambda_p$ and $1/\lambda_p$ (see figure 8). In the regime we consider they are all well below m_s. However, since the fundamental string tension is renormalized (82), in order for the mesons of section 4 to be well separated from the string excitations, they must be much lighter than the local string scale at $u \simeq \xi$. One can check that for the light mesons (*i.e.* those with $m \sim \sqrt{q}/\lambda_p$) this follows from (22), while for the heavy ones (those with $m \sim 1/\lambda_p$) it leads to the requirement

$$q \gg \left(\frac{\lambda_p}{\lambda_N}\right)^{\frac{2}{3}} = \left(\frac{p}{N}\right)^{\frac{2}{3}}. \tag{85}$$

This condition is stronger, but it still allows q to be very small.

6 Finite temperature

Holography relates the $4+1$ dimensional theory of N $D4$-branes at finite temperature to string theory in the near-horizon geometry of Euclidean non-extremal fourbranes [25]. This geometry is given by[17]

$$\begin{aligned} ds^2 &= H^{-1/3}[fd\tau^2 + dx_i^2 + dx_6^2 + dx_{11}^2] + H^{2/3}[(f^{-1}-1)dr^2 + |dv|^2 + |dw|^2 + dx_7^2], \\ C_6 &= H^{-1} d^4x \wedge dx_6 \wedge dx_{11}. \end{aligned} \tag{86}$$

Here we define $r^2 = |v|^2 + |w|^2 + x_7^2$, which obeys $r = u$ at $x_7 = 0$ (where all our branes from here on will be localized). In addition to the harmonic function H defined in (14), which in this limit takes the form

$$H = \frac{u_0^3}{r^3}, \qquad u_0^3 \equiv \pi \lambda_N l_s^2, \tag{87}$$

this metric contains a non–extremality factor f, which depends on the temperature T,

$$f = 1 - \frac{u_T^3}{r^3}, \qquad u_T = \frac{16\pi^2}{9}(Tl_s)^2 \lambda_N. \tag{88}$$

The Euclidean time $\tau = it$, is periodically identified,

$$\tau \sim \tau + \frac{1}{T}. \tag{89}$$

The relation between u_T and the temperature ensures the smoothness of the metric (86) at $u = u_T$.

To study the system of section 2 at finite temperature we need to place the NS, NS' and p $D4$-branes in the background (86). Since these branes are treated as probes, they do not change the background. One can think of (86) as providing a thermal bath in which the four dimensional confining gauge theory is placed. Even though the bath is five dimensional while the theory we are interested in is four dimensional, it is clear that

---

[17] As before, the type IIA geometry is obtained by reducing on the x_{11} circle.

T above is also the temperature felt by the four dimensional degrees of freedom, which are in thermal equilibrium with the five dimensional ones.[18]

Since the geometry is modified by the temperature, we need to determine the shape of the curved fivebrane in the new geometry. The free energy is given (in the probe approximation) by the fivebrane action for this shape (times the temperature). Recall that at zero temperature, the profile (6) has a $U(1)$ symmetry, (9), which corresponds to a $U(1)$ global symmetry in the gauge theory of section 3. From the field theory point of view, it is clear that this symmetry remains unbroken at finite temperature. In the geometry (86) this is the statement that the function f is invariant under (9). The most general ansatz consistent with this $U(1)$ is given by (15). Plugging it into the fivebrane Lagrangian (for $x_7 = 0$) leads to

$$L = H^{-1}\sqrt{f(1+H(u\dot{\phi})^2)\left(1+\frac{H}{f}(u')^2+H(u\alpha')^2\right)} - H^{-1}, \tag{90}$$

which is the finite temperature counterpart of the Lagrangian (17). As in the zero temperature case, the equations of motion imply that $\phi = x_{11}/\lambda_p$, and translation invariance in α and x_6 leads to the conserved charges

$$\begin{aligned} J &= \frac{u^2\alpha'\sqrt{f(1+Hu^2/\lambda_p^2)}}{\sqrt{1+\frac{H}{f}(u')^2+H(u\alpha')^2}} , \\ E &= H^{-1} - \frac{H^{-1}\sqrt{f(1+Hu^2/\lambda_p^2)}}{\sqrt{1+\frac{H}{f}(u')^2+H(u\alpha')^2}} . \end{aligned} \tag{91}$$

To study the branes at finite temperature it is convenient to invert the relation $u(x_6)$ and use u, rather than x_6, as an independent variable. After some algebra, equation (91) can be rewritten as

$$\frac{d\alpha}{du} = \frac{J}{u^2}\frac{dx_6}{du}\frac{1}{1-EH} , \tag{92}$$

$$\frac{dx_6}{du} = \pm\frac{u}{J\sqrt{f}}(1-EH)\left[\frac{u^2}{J^2}f(H^{-1}+\frac{u^2}{\lambda_p^2})-1-\frac{u^2}{J^2}(1-EH)^2H^{-1}\right]^{-1/2} . \tag{93}$$

It is convenient to define

$$G(u) = \frac{u^2}{J^2}f(H^{-1}+\frac{u^2}{\lambda_p^2})-1-\frac{u^2}{J^2}(1-EH)^2H^{-1} , \tag{94}$$

in terms of which (92), (93) take the form

$$\frac{d\alpha}{du} = \pm\frac{1}{u\sqrt{fG}} , \tag{95}$$

$$\frac{dx_6}{du} = \pm\frac{u}{J\sqrt{fG}}(1-EH) . \tag{96}$$

---

[18] Similar issues arise in the Sakai-Sugimoto model and related models, see, e.g., [26, 27, 28].

Note that x_6 is not a single–valued function of u, and we have to consider two branches, which correspond to the two signs in (95), (96). These branches connect at the point where u reaches its minimal value u_{min}, which is determined by the condition

$$G(u_{min}) = 0. \tag{97}$$

The integrals of motion, E and J, are determined by imposing appropriate boundary conditions on x_6 and α. As discussed in section 2.1, at large u the distance between the $NS5$-branes goes to infinity, so to define the theory one has to introduce a cutoff u_∞ and to impose boundary conditions at $u = u_\infty$.

The boundary has two components, corresponding to the NS and NS'-branes (or, equivalently, negative and positive branches of (95), (96)). On the negative (NS) branch, the boundary conditions are $\alpha(u_\infty) = \epsilon$, $x_6(u_\infty) = -L/2$; on the positive one they are $\alpha(u_\infty) = \pi/2 - \epsilon$, $x_6(u_\infty) = L/2$, where $\epsilon \to 0$ as $u_\infty \to \infty$. The boundary conditions preserve the symmetry of the equations of motion $x_6 \to -x_6$; thus, $\alpha(u_{min}) = \pi/4$.

It is important to emphasize that u_∞, L, ϵ above are independent of temperature – the boundary conditions are used to define the theory at the UV cutoff scale, and are independent of the state.[19] In particular, ϵ can be calculated in the zero temperature theory by using (12).

The constants J and E (91) can be calculated as a function of temperature by integrating the equations of motion. Consider first the integral of (95). In principle we should integrate from $u_{\rm min}$ to u_∞, but the resulting integral is convergent, so one can send the upper limit of integration to infinity (and thus $\epsilon \to 0$). This leads to

$$\int_{u_{min}}^{\infty} \frac{du}{u\sqrt{fG}} = \frac{\pi}{4}. \tag{98}$$

This, together with (97), gives one condition on J, E.

The second condition comes from integrating (96). This integral is divergent at large u, so we need to be more careful with it. Recall that at zero temperature the theory is characterized by the "QCD scale" ξ, which enters the relation (10) between L and u_∞. Denoting the integration constant related to ξ via (20) by J_0, and the corresponding function G (94) by G_0,[20] we can write (10) as

$$\frac{L}{2} = \int_{\sqrt{J_0\lambda_p}}^{u_\infty} \frac{udu}{J_0\sqrt{G_0}}. \tag{99}$$

Integrating the finite temperature equation of motion (96) leads to the relation

$$\frac{L}{2} = \int_{u_{min}}^{u_\infty} \frac{udu}{J\sqrt{fG}}(1 - EH) = \int_{\sqrt{J_0\lambda_p}}^{u_\infty} \frac{udu}{J_0\sqrt{G_0}}. \tag{100}$$

---

[19] Of course, we must choose the UV cutoff to be sufficiently large, $u_\infty \gg u_T$.

[20] So $G_0 = (u^2/J_0\lambda_p)^2 - 1$.

To remove the UV cutoff, it is useful to rewrite (100) such that the limit $u_\infty \to \infty$ is smooth. This can be done by subtracting the two integrals in (100) and combining them. The resulting integral is finite in the limit $u_\infty \to \infty$. In this limit one finds an integral equation in which u_∞ and L have been traded for the physical ("QCD") scale ξ. This is the brane analog of the process of renormalization in QCD.

Equations (97), (98) and (100) determine E and J as functions of J_0 (or ξ) and the temperature. The profile of the brane is then determined by solving (95) and (96). The free energy of the solution (divided by the temperature) is given by

$$\begin{aligned}\mathcal{E}^{(con)} &= \int_{u_{min}}^{u_\infty} du H^{-1}\left[\sqrt{f(1+\frac{u^2 H}{\lambda_p^2})\left((\partial_u x_6)^2+\frac{H}{f}+H(u\partial_u\alpha)^2\right)}-H^{-1}\partial_u x_6\right]\\ &= \int_{u_{min}}^{u_\infty}\frac{udu}{\sqrt{fJ^2G}}\left[H^{-1}(f-1)+\frac{u^2}{\lambda_p^2}f+E\right], \qquad (101)\end{aligned}$$

where we restricted to the positive branch; therefore the full free energy of the curved fivebrane is $2\mathcal{E}^{(con)}$.

The connected solution which has been discussed so far, corresponds to a confining vacuum of the gauge theory of section 3 at finite temperature. Another solution of the equations of motion, which corresponds to the Higgs branch, is a configuration of two disconnected fivebranes which descend from large values of u to the horizon located at $u = u_T$. Looking at equation (93) and requiring x_6' to be real for all $u > u_T$, we find that such a disconnected solution must have

$$J = 0, \qquad E = \frac{u_T^3}{u_0^3}. \qquad (102)$$

For these values of parameters equation (96) simplifies (we only look at one of the branches)

$$\frac{dx_6}{du} = \left(\frac{u^2}{\lambda_p^2}+\frac{u_T^3}{u_0^3}\right)^{-\frac{1}{2}}. \qquad (103)$$

It has a unique solution satisfying the relation (99) between L and u_∞,

$$x_6 = \lambda_p \ln\left(u+\sqrt{u^2+\frac{\lambda_p^2 u_T^3}{u_0^3}}\right)-\frac{\lambda_p}{2}\ln(2J_0\lambda_p). \qquad (104)$$

This solution should only be considered in the exterior of the black hole, $u > u_T$. The derivative $\frac{d\alpha}{du}$ vanishes in this region (see (92)); the solution has $\alpha = \pi/2$ and describes an NS'-brane with p $D4$-branes attached.

At zero temperature the connected and disconnected solutions have the same energy (in the limit when the cutoff u_∞ is sent to infinity), in agreement with the fact that they describe two supersymmetric vacua of the same theory. At finite temperature, one of them can have lower free energy. The free energy of the connected solution is given by twice (101); for the disconnected one we find a free energy $2\mathcal{E}^{(dis)}$ with

$$\begin{aligned}
\mathcal{E}^{(dis)} &= \int_{u_T}^{u_\infty} du \left[\sqrt{f(H^{-1}+\frac{u^2}{\lambda_p^2})\left(H^{-1}(\partial_u x_6)^2+\frac{1}{f}\right)} - H^{-1}\partial_u x_6\right] \\
&= \frac{1}{2}\left[u\sqrt{\frac{u^2}{\lambda_p^2}+\frac{u_T^3}{u_0^3}} - \lambda_p \frac{u_T^3}{u_0^3}\ln\left\{u+\sqrt{u^2+\lambda_p^2\frac{u_T^3}{u_0^3}}\right\}\right]_{u_T}^{u_\infty}. \qquad (105)
\end{aligned}$$

Let us demonstrate that at small but finite value of u_T this free energy is smaller than (101). Near $u_T = 0$, (105) behaves as

$$\mathcal{E}^{(dis)} = \frac{1}{2\lambda_p}\left[u_\infty^2 + \frac{\lambda_p^2 u_T^3}{2u_0^3}(1+\ln\left(\frac{u_T^2}{u_\infty^2}\right)) - u_T^2 + o(u_T^3)\right]. \qquad (106)$$

This should be compared with a similar expansion of (101),

$$\begin{aligned}
\mathcal{E}^{(con)} &= \int_{u_{min}}^{u_\infty} \frac{udu}{\sqrt{fJ^2G}}\left[\frac{u^2}{\lambda_p^2}+E\right] - u_T^3 \int_{u_{min}}^{u_\infty} \frac{udu}{u^3\sqrt{fJ^2G}}\left[H^{-1}+\frac{u^2}{\lambda_p^2}\right] \qquad (107) \\
&= \frac{1}{\lambda_p^2}\int_{u_{min}}^{u_\infty} \frac{u^3 du}{\sqrt{fJ^2G}} - u_T^3 \int_{\sqrt{J_0\lambda_p}}^{u_\infty} \frac{du}{u^2\sqrt{u^4/\lambda_p^2 - J_0^2}}\left[H^{-1}+\frac{u^2}{\lambda_p^2}-\frac{E}{u_T^3}u^3\right] + o(u_T^3).
\end{aligned}$$

Here we used the fact that the ratio E/u_T^3 remains finite as u_T goes to zero (this can be shown by performing a small u_T expansion of (98) and (100)) as well as (94). To simplify the first term in (107), we observe that

$$fJ^2G = \frac{u^4}{\lambda_p^2} - J^2 + 2Eu^2 + \frac{u_T^3}{u^3}\left[J^2 - H^{-1}u^2 - 2\frac{u^4}{\lambda_p^2}\right] + o(u_T^3). \qquad (108)$$

This equation leads to an expression for du^4 in terms of $d(fJ^2G)$, and substituting the result into the first term in (107), we find

$$\begin{aligned}
&\frac{1}{4\lambda_p^2}\int_{u_{min}}^{u_\infty} \frac{du^4}{\sqrt{fJ^2G}} \\
&\quad = \frac{1}{4}\left[2\sqrt{fJ^2G}\Big|_{u_{min}}^{u_\infty} - u_T^3 \int_{u_{min}}^{u_\infty} \frac{1}{\sqrt{fJ^2G}} d\left\{\frac{2E}{u_T^3}u^2 + \frac{J^2}{u^3} - \frac{1}{uH} - 2\frac{u}{\lambda_p^2}\right\} + o(u_T^3)\right] \\
&\quad \sim \frac{u_\infty^2}{4\lambda_p}\left(2 - \frac{\lambda_p^2}{u_\infty^2}\frac{u_T^3}{u_0^3} + 2E\frac{\lambda_p^2}{u_\infty^2}\right) \\
&\qquad - \frac{u_T^3}{4}\int_{\sqrt{J_0\lambda_p}}^{u_\infty} \frac{du}{\sqrt{u^4/\lambda_p^2 - J_0^2}}\left\{\frac{4E}{u_T^3}u - \frac{3J^2}{u^4} - \frac{2u}{u_0^3} - \frac{2}{\lambda_p^2}\right\} + o(u_T^3). \qquad (109)
\end{aligned}$$

At the last stage we dropped some terms which vanish in the limit $u_\infty \to \infty$. Substituting (109) into (107) and performing the integral in the term that diverges at large u_∞, we find

$$\begin{aligned}\mathcal{E}^{(con)} &= \frac{u_\infty^2}{2\lambda_p} + u_T^3\left[\frac{\lambda_p E}{2u_T^3} - \frac{\lambda_p}{4u_0^3}\right] + \frac{\lambda_p u_T^3}{4u_0^3}\ln\left(\frac{J_0\lambda_p}{u_\infty^2}\right) \\ &+ \frac{u_T^3}{4}\int_{\sqrt{J_0\lambda_p}}^{u_\infty}\frac{du}{\sqrt{u^4/\lambda_p^2 - J_0^2}}\left\{\frac{3J^2}{u^4} - \frac{2}{\lambda_p^2}\right\} + o(u_T^3)\end{aligned} \tag{110}$$

For small values of the nonextremality parameter u_T, this expression is larger than (106),

$$\mathcal{E}^{(con)} - \mathcal{E}^{(dis)} = \frac{u_T^2}{2\lambda_p} + \frac{\lambda_p u_T^3}{4u_0^3}\ln\left(\frac{J_0\lambda_p}{u_T^2}\right) + O(u_T^3). \tag{111}$$

Thus, we see that the disconnected configuration is thermodynamically preferred for all non-zero T, and the phase transition from the confining to the Higgs phase (if we start from the confining phase at zero temperature) occurs at $T = 0$. Of course, we can still study the confining phase at (sufficiently small) non-zero T, but it is meta-stable in this regime.

The above discussion is natural from the field theory point of view. The Higgs phase of the gauge theory discussed in section 3 (corresponding to the brane configuration of figure 4) has $(N+p)^2$ massless fields, while the confining phase has N^2. Since the former has more massless degrees of freedom, its free energy is lower; the difference is an order p/N effect. Our result on the energetics of the branes implies that this behavior persists at strong coupling.

It is natural to ask whether it is possible to deform the brane system so as to shift the deconfinement phase transition to finite temperature. One way to do that is to start at zero temperature with a brane configuration in which the $NS5$-branes are displaced relative to the N $D4$-branes in the x_7 direction by the distance $X_7 \neq 0$. The free energies of the confining (figure 2) and Higgs (figure 4) branches depend on X_7, and it is possible that the transition between them occurs at finite T.

A complication in this analysis is that at finite temperature the curved connected and disconnected fivebranes that correspond to the two branches are no longer located at fixed x_7. One can show that they develop a profile in x_7 that (at large x_6) changes logarithmically with u. This bending can be understood from field theory as due to the fact that in order to have a non-zero vacuum expectation value of x_7 at finite temperature, one has to add to the Lagrangian a $4+1$ dimensional FI D-term for the $U(1)$'s on the semi-infinite $D4$-branes. The presence of this logarithmic mode complicates the analysis of the energetics, and we will leave it to future work.

7 Non-supersymmetric generalizations

So far we have discussed brane configurations with $d = 4$ $\mathcal{N} = 1$ supersymmetry. There are many possible generalizations, to different dimensions and different amounts of supersymmetry. In this section we discuss two non-supersymmetric brane systems that are closely related to those studied in this paper. The first is a configuration similar to that discussed in section 2.2 and drawn in figure 3, but with the two $NS5$-branes taken to be parallel and having opposite orientations (so that in the classical limit they look like an NS-brane and an $\overline{NS}$-brane). We analyze this model both at zero and at finite temperature. The second system is similar to the model discussed in section 2.3, with a compact x_6 direction (see figure 5), but with anti-periodic boundary conditions for the fermions around the circle.

7.1 The $NS - \overline{NS}$ system

We start with the brane configuration of figure 3, with the NS'-brane replaced by a second $NS5$-brane parallel to the first one, but with opposite orientation; we will refer to it as an $\overline{NS}$-brane. In the full string theory, this configuration is unstable to gravitational attraction of the fivebranes, but in the decoupled theory of the N $D4$-branes, which we focus on here, the mode governing the separation between the fivebranes is non-normalizable, and this instability is absent.

The field theory in this case is similar to the one discussed in section 3 for the configuration of figure 3. In that case the couplings of the fundamental fields living at the brane intersections to the adjoint fields preserved $\mathcal{N} = 2$ supersymmetry at each intersection separately, but only $\mathcal{N} = 1$ for both intersections together, while in this case there are no common supersymmetries. Classically the brane configuration has a moduli space corresponding to moving the $D4$-branes on the interval along the $NS5$-branes; quantum mechanically, we expect this moduli space to be lifted. It would be interesting to see whether at weak coupling the field theory potential on this moduli space is attractive or repulsive.

In the string dual, this brane configuration (for $x_7 = 0$) is described by the Lagrangian (17) with $\alpha' = 0$. Expressing the Lagrangian in terms of dimensionless coordinates

$$u \to \frac{u}{\hat{u}}, \quad x_6 \to \frac{x_6}{\lambda_p}, \qquad \hat{u} \equiv \frac{\pi \lambda_N l_s^2}{\lambda_p^2}, \tag{112}$$

as was done above equation (49), we find (up to an overall constant)

$$\mathcal{L} = u^{5/2}\sqrt{(1+u)\left(1 + \frac{(u')^2}{u^3}\right)} - u^3. \tag{113}$$

The corresponding "conserved Hamiltonian" is given by

$$E = u^3 - \frac{u^{5/2}\sqrt{(1+u)}}{\sqrt{1 + \frac{(u')^2}{u^3}}}. \tag{114}$$

This implies the following equation

$$u' = \pm u^{3/2}\sqrt{\frac{u^5(1+u)}{(u^3-E)^2} - 1}. \tag{115}$$

It is easy to see that for $E = 0$ there is no connected solution but only a disconnected one, of the form (21) $u = K\exp(\pm x_6/\lambda_p)$, describing a configuration in which the gauge symmetry is broken as in section 3. In the discussion of section 2, the connected solution also had $E = 0$ due to supersymmetry, but now with no supersymmetry there is no reason why a connected solution should have this property. Connected solutions exist for any $E < 0$, and they are characterized by having a minimum point where $u' = 0$. This point is the maximal solution $u = u_*$ to the equation for the vanishing of u',

$$u_*^5 + 2Eu_*^3 - E^2 = 0. \tag{116}$$

It is easy to check (this is a special case of a computation done in the next subsection) that the connected configuration always has lower energy than the disconnected one, so it describes the vacuum of this non-supersymmetric field theory. We did not analyse explicitly the stability of this brane configuration, but we expect it to be stable, and to otherwise exhibit similar physics to what we found in the previous sections.

The theory is parametrized by u_* (the minimal value of u), which is an analog of the "QCD scale" ξ introduced in section 2. Integrating (115) between u_* and the UV cutoff u_∞, we find an analog of the relation (10) between the QCD scale and the distance between the branes at $u = u_\infty$:

$$\frac{L}{2} = \int_{u_*}^{u_\infty} \frac{(u^3-E)du}{\sqrt{u^3(u^5+2Eu^3-E^2)}}. \tag{117}$$

Here $E(u_*)$ is a negative number that solves equation (116). The integral on the right–hand side of (117) diverges logarithmically for large u_∞, so it is convenient to define a renormalized quantity,

$$L_{ren} = \lim_{u_\infty\to\infty}(L - 2\ln(u_\infty)) = 2\int_{u_*}^{\infty}\left[\frac{(u^3-E)}{\sqrt{u^3(u^5+2Eu^3-E^2)}} - \frac{1}{u}\right]du - 2\ln(u_*), \tag{118}$$

which remains finite as $u_\infty \to \infty$. Inverting this relation, one finds u_* as a function of L_{ren}, which is shown in the red line in figure 15.

7.2 The $NS - \overline{NS}$ system at finite temperature

At finite temperature, the $NS - \overline{NS}$ system is described by the Lagrangian (90) with $\alpha' = 0$. Performing the re-scaling (112) in (90), or equivalently introducing the thermal factor $f(u) = 1 - u_T^3/u^3$ into the Lagrangian (113), we find[21]

$$\mathcal{L} = u^{5/2}\sqrt{f(1+u)\left(1+\frac{(u')^2}{fu^3}\right)} - u^3. \tag{119}$$

The corresponding Hamiltonian for translations in x_6 is given by

$$E_T = u^3 - \frac{u^{5/2}\sqrt{f(1+u)}}{\sqrt{1+\frac{(u')^2}{fu^3}}}. \tag{120}$$

The minimal value of u for the connected configuration, $u = u_{min}$, corresponds to $u' = 0$, which is given by the solution of

$$u_{min}^5 - u_T^3(u_{min}^2 + u_{min}^3) + 2E_T u_{min}^3 - E_T^2 = 0. \tag{121}$$

The disconnected configuration has $u' = 0$ at $u_{min} = u_T$, where according to (121) $E = u_T^3$.

The connected configuration is characterized by the asymptotic separation L, which is given in terms of u_{min} and u_T as follows

$$L = 2\int_{u_{min}}^{u_\infty} \frac{du}{u'} = 2\int_{u_{min}}^{u_\infty} \frac{u^3 - E_T}{u^{3/2}\sqrt{f}} \frac{1}{\sqrt{u^5 - u_T^3(u^2+u^3) + 2E_T u^3 - E_T^2}}\,du. \tag{122}$$

The separation L can be renormalized in the same manner as in (118). The action of the connected configuration is given by (up to an overall constant)

$$S^{(con)} = 2\int_{u_{min}}^{u_\infty} \frac{u^{3/2}}{\sqrt{f}} \frac{f(u^3+u^2) - u^3 + E_T}{\sqrt{u^5 - u_T^3(u^2+u^3) + 2E_T u^3 - E_T^2}}\,du\,, \tag{123}$$

while the action of the disconnected configuration is

$$S^{(dis)} = 2\int_{u_T}^{u_\infty} \frac{u^2 du}{\sqrt{u^2 + u_T^3}}\,. \tag{124}$$

The last integral can be evaluated exactly leading to

$$S^{(dis)} = \left[u\sqrt{u^2+u_T^3} - u_T^3 \ln[2(u + \sqrt{u^2+u_T^3})]\right]_{u_T}^{u_\infty}\,. \tag{125}$$

The question of which of the two configurations, the connected one or the disconnected one, is preferable at any given temperature is determined by the difference of the corresponding free energies. In our approximation this is translated into the difference between

---

[21] Notice that this Lagrangian differs from (90) by an overall factor, and that u_T here is rescaled with respect to u_T of section 6.



Figure 13: ΔS as a function of L_{ren} for $u_T = 0.5$ (blue), $u_T = 1$ (red), and $u_T = 1.5$ (yellow). The arrows indicate in which direction the position of the brane u_{min} increases.

the classical actions of the connected and disconnected configurations. Each action separately is divergent as $u_\infty \to \infty$, but the difference between them is finite, and is given by

$$\begin{aligned}\Delta S \equiv S^{(con)} - S^{(dis)} &= 2\int_{u_{min}}^{\infty} u^{3/2}\left\{\frac{1}{\sqrt{f}}\frac{fu^2 - u_T^3 + E_T}{\sqrt{u^5 - u_T^3(u^2+u^3) + 2E_T u^3 - E_T^2}} - \frac{u^2}{\sqrt{u^2+u_T^3}}\right\} du \\ &\quad - \left[u\sqrt{u^2+u_T^3} - u_T^3 \ln[2(u+\sqrt{u^2+u_T^3})]\right]_{u_T}^{u_{min}} . \end{aligned} \tag{126}$$

The difference of the actions as a function of the asymptotic separation distance is shown, for some specific values of the temperature, in figure 13.

The qualitative dependence of ΔS on L depicted in the figures is the same for any temperature $u_T > 0$. From the figures it is evident that ΔS is always negative, and hence the connected solution is preferred. However, it is also evident that for any temperature there is a critical asymptotic separation distance L_{cr} (for a fixed UV cutoff), above which the connected configuration no longer exists. This is manifest in figure 14 which describes the separation distance L as a function of u_{min} for a particular temperature. Increasing u_{min} from u_T and correspondingly changing E_T according to (121), L first increases to L_{cr}, and then decreases. Therefore, for any given separation distance L, when the temperature is raised, eventually L becomes larger than the critical value and hence the connected configuration will cease to exist. The dependence of the phase transition temperature u_T on the renormalized L_{cr}, together with the corresponding value of u_* at $T = 0$, are shown in figure 15.



Figure 14: The asymptotic separation distance L_{ren} as a function of u_{min} for $u_T = 1$.



Figure 15: u_* (red,higher) (at $T = 0$), and the phase transition temperature u_T (blue,lower) as a function of the renormalized distance L_{ren}.

The precise value of the phase transition point may be found by solving $\partial L/\partial u_{min} = 0$ for u_{min} (at fixed temperature u_T), with $L(u_{min})$ given by (122); note that this depends on u_{min} both through E_T and through the lower bound on the integration region (and that the resulting equation has a finite limit as $u_\infty \to \infty$). Inverting the function $u_{min}(u_T)$ that we find gives us the phase transition temperature for a given value of u_{min} and of $L = L(u_{min})$. At this temperature there will be a first order phase transition to a disconnected configuration. We have thus shown that unlike the supersymmetric case discussed in section 6, for the non-supersymmetric model there is a first order phase transition between the two phases at non-zero temperature.

7.3 A fivebrane in the cigar topology

The second non-supersymmetric model we discuss is similar to the compactified model of section 2.3, but with anti-periodic boundary conditions for the fermions. This system, which breaks supersymmetry, is not stable in flat space, but it is stable in the near-horizon limit of the $D4$-branes [25]. The bulk geometry in this case is a double-Wick-rotation of the near-extremal $D4$-brane solution, and in this background the (x_6, u) coordinates have the topology of a cigar (as opposed to the cylinder discussed in section 2.3). The eleven dimensional background is given by (compare to (86))

$$\begin{aligned} ds^2 &= H^{-1/3}[-dt^2 + dx_i^2 + f(r)dx_6^2 + dx_{11}^2] + H^{2/3}[(f(r)^{-1} - 1)dr^2 + |dv|^2 + |dw|^2 + dx_7^2], \\ C_6 &= H^{-1} d^4x \wedge dx_6 \wedge dx_{11}. \end{aligned} \tag{127}$$

The function f is given by $f(r) = 1 - \left(\frac{u_\Lambda}{r}\right)^3$, where u_Λ is the value of the radial coordinate r at the tip of the cigar.

Upon reduction to ten dimensional type IIA supergravity, the corresponding metric, dilaton and RR form read

$$\begin{aligned} ds^2 &= \left(\frac{r}{R_{D4}}\right)^{3/2}\left[-dt^2 + \delta_{ij}dx^i dx^j + f(r)dx_6^2\right] + \left(\frac{R_{D4}}{r}\right)^{3/2}\left[\frac{dr^2}{f(r)} + r^2 d\Omega_4^2\right], \\ F_4 &= \frac{2\pi N_c}{V_4}\epsilon_4 \;\;, \;\; e^\phi = g_s\left(\frac{r}{R_{D4}}\right)^{3/4}, \;\; R_{D4}^3 = u_0^3 = \pi g_s N_c l_s^3. \end{aligned} \tag{128}$$

Note that the transformation from the extremal background to the non-extremal one is not just a change of the warp factor $H(r)$ which, as we have seen in (19), does not modify the brane profile. Fermions in the cigar background must be taken to be anti-periodic around the x_6 circle. Thus, supersymmetry is broken.

The fivebrane Lagrangian (for $x_7 = 0$) that follows from (128), again rescaling (112), is given by (up to a multiplicative constant)

$$\mathcal{L} = u^{5/2}\sqrt{1+u}\sqrt{\left(1 - \frac{u_\Lambda^3}{u^3}\right) + \frac{(u')^2 + (u\alpha')^2}{u^3 - u_\Lambda^3}} - u^3, \tag{129}$$



Figure 16: The spiraling profile over the cigar background. Again, we used a red line for the “downward” brane and a blue one for the climbing one.

The Hamiltonian for translations in x_6 is now

$$E = u^3 - \frac{u^{5/2}(1-\frac{u_\Lambda^3}{u^3})\sqrt{1+u}}{\sqrt{(1-\frac{u_\Lambda^3}{u^3}) + \frac{(u')^2+(u\alpha')^2}{u^3-u_\Lambda^3}}}\,. \qquad (130)$$

The conserved charge associated with the shift symmetry of α is

$$J = \frac{u^{5/2}\sqrt{1+u}\left(\frac{\alpha'}{u\left(1-\frac{u_\Lambda^3}{u^3}\right)}\right)}{\sqrt{(1-\frac{u_\Lambda^3}{u^3}) + \frac{(u')^2+(u\alpha')^2}{u^3-u_\Lambda^3}}}\,. \qquad (131)$$

In general it is hard to solve these equations exactly; it is simpler to do this in the case $E = 0$, but this case now does not lead to the correct change in α to connect the NS and NS'-branes. We expect that there should still exist an infinite sequence of connected solutions, as discussed in section 2.3 and as depicted in figure 16, but now these solutions will no longer be degenerate. It would be interesting to analyze these solutions in detail, to see which solution has the lowest energy. In the limit $\xi \gg u_\Lambda$ we can analyze this by working at leading order in u_Λ; we find that when the brane lies in a region that the parameter q defined in (49) obeys $q \ll 1$, the solutions with lower ξ (more cascade steps) have lower energy, so that the minimal energy configuration has ξ close to u_Λ, while for $q \gg 1$ the situation is the opposite. Due to the topology of the background it is obvious that in the present case we cannot have a disconnected profile, since the $NS5$-brane does not have where to end.

Acknowledgements

We thank A. Giveon for discussions. The work of OA, DK, JS and SY was supported in part by the Israel-U.S. Binational Science Foundation. The work of OA, JS and SY was supported in part by a research center supported by the Israel Science Foundation (grant number 1468/06) and by a grant (DIP H52) of the German Israel Project Cooperation. The work of OA was supported in part by the Minerva foundation with funding from the Federal German Ministry for Education and Research. The work of DK was supported in part by DOE grant DE-FG02-90ER40560 and NSF grant 0529954. The work of OL was supported in part by NSF grant 0844614. DK thanks the Weizmann Institute and OL thanks the University of Chicago for hospitality during part of this work.

A Approximations in the derivation of the spectra of the fluctuations in the transverse directions

The equations of motion that follow from the Lagrangian (73) are rather complicated, so to get at least some qualitative understanding on the structure of the bound states we invoke several approximations. Since the dominant contribution of $\frac{H_0}{((pR)^2+u^2H)_0}$ comes from the vicinity of $x_6 = 0$, where $V_0 \sim W_0$, to get a rough estimate of the eigenvalues we replace V_0/u_0 and W_0/u_0 by $\frac{1}{\sqrt{2}}$, and hence

$$[V_0V_1' - W_0W_1' - V_0V_1 - W_0W_1]^2 \approx \frac{u_0^2}{2}[V_1' - W_1' - V_1 - W_1]^2 = u_0^2\left[V_-' - V_+\right]^2 . \qquad (132)$$

Here we introduced

$$V_+ = \frac{V_1 + W_1}{\sqrt{2}}, \qquad V_- = \frac{V_1 - W_1}{\sqrt{2}}. \qquad (133)$$

This approximation leads to the Lagrangian

$$L \approx \frac{1}{2}\int dx_6 \left[(V_+')^2 + (V_-')^2 + \left(1 - (mpR)^2 - \frac{(mpR)^2}{q}\frac{1}{\sqrt{2\cosh[2x_6]}}\right)(V_+^2 + V_-^2)\right.$$

$$
-\frac{(mpR)^2}{q}\frac{1}{\sqrt{2\cosh[2x_6]}}\left[(mpR)^2+\frac{(mpR)^2}{q}\frac{1}{\sqrt{2\cosh[2x_6]}}\right]^{-1}(V_+-V_-')^2\Bigg]
$$

$$
\equiv\ \frac{1}{2}\int dx_6\left[(V_+')^2+(V_-')^2+a(x_6)(V_+^2+V_-^2)+b(x_6)(V_+-V_-')^2\right]. \qquad (134)
$$

This is still a complicated coupled Lagrangian. In the limit of $q << 1$ we may be justified to assume

$$
-a\sim\frac{(mpR)^2}{q}\frac{1}{\sqrt{2\cosh[2x_6]}}\gg 1, \qquad b\approx 1, \qquad b'\sim q\ll 1. \qquad (135)
$$

Using this approximation we get a decoupling of V_+ and V_- in the Lagrangian so that for the former we find

$$
L\ \approx\ \frac{1}{2}\int dx_6\left[(V_+')^2+\left(1-(mpR)^2-\frac{(mpR)^2}{q}\frac{1}{\sqrt{2\cosh[2x_6]}}\right)V_+^2\right]. \qquad (136)
$$

References

[1] G. 't Hooft, "A planar diagram theory for strong interactions," Nucl. Phys. B **72**, 461 (1974).

[2] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, "Large N field theories, string theory and gravity," Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111].

[3] C. Nunez, A. Paredes and A. V. Ramallo, "Unquenched flavor in the gauge/gravity correspondence," arXiv:1002.1088 [hep-th].

[4] A. Giveon and D. Kutasov, "Brane dynamics and gauge theory," Rev. Mod. Phys. **71**, 983 (1999) [arXiv:hep-th/9802067].

[5] J. Polchinski and M. J. Strassler, "The string dual of a confining four-dimensional gauge theory," arXiv:hep-th/0003136.

[6] I. R. Klebanov and M. J. Strassler, "Supergravity and a confining gauge theory: Duality cascades and chiSB-resolution of naked singularities," JHEP **0008**, 052 (2000) [arXiv:hep-th/0007191].

[7] J. M. Maldacena and C. Nunez, "Towards the large N limit of pure N = 1 super Yang Mills," Phys. Rev. Lett. **86**, 588 (2001) [arXiv:hep-th/0008001].

[8] N. Seiberg, "Electric - magnetic duality in supersymmetric nonAbelian gauge theories," Nucl. Phys. B **435**, 129 (1995) [arXiv:hep-th/9411149].

[9] E. Witten, "Branes and the dynamics of QCD," Nucl. Phys. B **507**, 658 (1997) [arXiv:hep-th/9706109].

[10] N. Itzhaki, J. M. Maldacena, J. Sonnenschein and S. Yankielowicz, "Supergravity and the large N limit of theories with sixteen supercharges," Phys. Rev. D **58**, 046004 (1998) [arXiv:hep-th/9802042].

[11] A. Dymarsky, I. R. Klebanov and N. Seiberg, "On the moduli space of the cascading $SU(M+p) \times SU(p)$ gauge theory," JHEP **0601**, 155 (2006) [arXiv:hep-th/0511254].

[12] E. Witten, "Solutions of four-dimensional field theories via M-theory," Nucl. Phys. B **500**, 3 (1997) [arXiv:hep-th/9703166].

[13] F. Benini, F. Canoura, S. Cremonesi, C. Nunez and A. V. Ramallo, "Backreacting Flavors in the Klebanov-Strassler Background," JHEP **0709**, 109 (2007) [arXiv:0706.1238 [hep-th]].

[14] O. Aharony and A. Hanany, "Branes, superpotentials and superconformal fixed points," Nucl. Phys. B **504**, 239 (1997) [arXiv:hep-th/9704170].

[15] I. R. Klebanov and E. Witten, "Superconformal field theory on threebranes at a Calabi-Yau singularity," Nucl. Phys. B **536**, 199 (1998) [arXiv:hep-th/9807080].

[16] I. R. Klebanov and N. A. Nekrasov, "Gravity duals of fractional branes and logarithmic RG flow," Nucl. Phys. B **574**, 263 (2000) [arXiv:hep-th/9911096].

[17] I. R. Klebanov and A. A. Tseytlin, "Gravity Duals of Supersymmetric $SU(N) \times SU(N+M)$ Gauge Theories," Nucl. Phys. B **578**, 123 (2000) [arXiv:hep-th/0002159].

[18] M. J. Strassler, "The duality cascade," arXiv:hep-th/0505153.

[19] O. Aharony, A. Giveon and D. Kutasov, "LSZ in LST," Nucl. Phys. B **691**, 3 (2004) [arXiv:hep-th/0404016].

[20] T. Sakai and S. Sugimoto, "Low energy hadron physics in holographic QCD," Prog. Theor. Phys. **113**, 843 (2005) [arXiv:hep-th/0412141].

[21] E. Antonyan, J. A. Harvey, S. Jensen and D. Kutasov, "NJL and QCD from string theory," arXiv:hep-th/0604017.

[22] M. Van Raamsdonk and K. Whyte, "Baryons from embedding topology and a continuous meson spectrum in a new holographic gauge theory," arXiv:0912.0752 [hep-th].

[23] P. Pasti, D. P. Sorokin and M. Tonin, "Covariant action for a D = 11 five-brane with the chiral field," Phys. Lett. B **398** (1997) 41 [arXiv:hep-th/9701037].

[24] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, “Wilson loops, confinement, and phase transitions in large N gauge theories from supergravity,” JHEP **9806**, 001 (1998) [arXiv:hep-th/9803263].

[25] E. Witten, “Anti-de Sitter space, thermal phase transition, and confinement in gauge theories,” Adv. Theor. Math. Phys. **2**, 505 (1998) [arXiv:hep-th/9803131].

[26] O. Aharony, J. Sonnenschein and S. Yankielowicz, “A holographic model of deconfinement and chiral symmetry restoration,” Annals Phys. **322**, 1420 (2007) [arXiv:hep-th/0604161].

[27] A. Parnachev and D. A. Sahakyan, “Chiral phase transition from string theory,” Phys. Rev. Lett. **97**, 111601 (2006) [arXiv:hep-th/0604173].

[28] E. Antonyan, J. A. Harvey and D. Kutasov, “The Gross-Neveu model from string theory,” Nucl. Phys. B **776**, 93 (2007) [arXiv:hep-th/0608149].